


01/13/00



jc658 U.S. PTO

Attorney Docket No.: 5778.200-US

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
FILING UNDER 37 C.F.R. 1.53(b)Box Patent Application
Assistant Commissioner for Patents
Washington, DC 20231Express Mail Label No.: EL443984197US
Date of Deposit: January 13, 2000jc678 U.S. PTO
09/482788

Sir:

This is a request for filing a **continuation-in-part** application under 37 C.F.R.

1.53(b) of

Applicant(s): Berka *et al.*Title: Methods For Producing Polypeptides In Cyclohexadepsipeptide-Deficient
Cells51 pages of specification 10 sheets of formal drawings 11 pages of Sequence Listing
3 sheets of Declaration and Power of Attorney☒ The filing fee is calculated as follows:

Basic Fee:	\$690.00
Total Claims: $20 - 20 = 0 \times 18 =$	\$0
Independent Claims: $3 - 3 = 0 \times 78 =$	\$0
Total Fee:	\$690.00

This application is a continuation-in-part of application no. 09/229,862 filed on
January 13, 1999 and claims priority under 35 U.S.C. 119, the contents of which are fully
incorporated herein by reference.Address all future communications to Steve T. Zelson, Esq., Novo Nordisk of North
America, Inc., 405 Lexington Avenue, Suite 6400, New York, NY 10174-6401.Please charge the required fee, estimated to be \$690, to Novo Nordisk of North
America, Inc., Deposit Account No. 14-1447. A duplicate of this sheet is enclosed.

Date: January 13, 2000

Respectfully submitted,

Robert L. Starnes, Reg. No. 41,324
Novo Nordisk of North America, Inc.
405 Lexington Avenue, Suite 6400
New York, New York 10017

09/482788-011300

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

EXPRESS MAIL CERTIFICATE

Box Patent Application
Assistant Commissioner for Patents
Washington, DC 20231

Re: U.S. Patent Application for
"Methods For Producing Polypeptides In Cyclohexadepsipeptide-
Deficient Cells"
Applicants: Berka *et al.*

Sir:

Express Mail Label No.: EL443984197US

Date of Deposit: January 13, 2000

I hereby certify that the following attached paper(s) or fee

1. Filing Under 37 C.F.R. 1.53(b) (in duplicate)
2. Patent Application
3. Executed Combined Declaration and Power of Attorney
4. Preliminary Amendment
5. Assignment
6. Recordation
7. Sequence Listing
8. Sequence Listing Diskette
9. Verified Statement

are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 C.F.R. 1.10 on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, DC 20231.

Selest A. Buriani

(Name of person mailing papers or fees)

Selest A. Buriani

(Signature of person mailing papers or fees)

Mailing Address:

Novo Nordisk of North America, Inc.
405 Lexington Avenue, Suite 6400
New York, NY 10017
(212) 867-0123

00110-9828-90

Please amend claim 63 as follows:

At line 1, delete "any of claims 30-52" and insert --claim 30--.

Please amend claim 64 as follows:


At line 1, delete "any of claims 30-52" and insert --claim 30--.

REMARKS

This amendment is submitted to cancel claims and correct improper multiple dependent claims. Since only claims are cancelled and dependencies are altered, there is no new matter added, and entry of the amendment is respectfully requested.

Respectfully submitted,

Date: January 13, 2000



Robert L. Starnes, Reg. No. 41,324
Novo Nordisk of North America, Inc.
405 Lexington Avenue, Suite 6400
New York, NY 10174-6401
(212) 867-0123

METHODS FOR PRODUCING POLYPEPTIDES IN CYCLOHEXADEPSIPEPTIDE-DEFICIENT CELLS

Cross-Reference to Related Application

This application is a continuation-in-part of pending U.S. application Serial No. 09/229,862 filed January 13, 1999, which application is fully incorporated herein by reference.

Background of the Invention

Field of the Invention

The present invention relates to methods for producing heterologous polypeptides in cyclohexadepsipeptide-deficient filamentous fungal mutant cells. The present invention also relates to mutants of filamentous fungal cells and methods for obtaining the mutant cells. The present invention also relates to isolated cyclohexadepsipeptide synthetases and isolated nucleic acid sequences encoding the cyclohexadepsipeptide synthetases. The present invention also relates to nucleic acid constructs, vectors, and host cells comprising the nucleic acid sequences as well as methods for producing the cyclohexadepsipeptide synthetases. The present invention further relates to cyclohexadepsipeptides produced by the cyclohexadepsipeptide synthetases.

Description of the Related Art

Depsipeptides constitute a large class of peptide-related compounds derived from hydroxy and amino acids joined by amide and ester linkages. Many members of this class of compounds are biologically active and include antibiotics, alkaloids, and proteins (Shemyakin *et al.*, 1969, *Journal of Membrane Biology* 1: 402-430). Examples include the enniatins, beauvericin, and bassianolide.

Enniatins are cyclohexadepsipeptide phyto toxins with ionophoretic properties produced by various species of actinomycetes and filamentous fungi, particularly strains of *Fusarium*. They are composed of alternating D-2-hydroxyisovaleric acid residues and L-amino acids or

N-methyl-L-amino acids to form an 18-membered cyclic structure and may contain more than one species of amino acid.

The biosynthesis of enniatins is catalyzed by enniatin synthetase, which is a large multifunctional enzyme that has all the essential functions for assembling enniatins from their primary precursors, *i.e.*, D-2-hydroxyisovaleric acid, a branched chain L-amino acid (*e.g.*, valine, leucine, isoleucine), S-adenosylmethionine, and ATP (Reper *et al.*, 1995, *European Journal of Biochemistry* 230: 119-126). The precursors (D-2-hydroxyisovaleric acid and branched chain L-amino acid) are activated as thioesters. Covalently bound substrate amino acid residues are methylated under the consumption of S-adenosylmethionine. Then peptide bond formation and cyclization reactions occur.

Enniatins are postulated to play a role in wilt toxic events during infection by enniatin-producing fusaria (Walton, 1990, *Biochemistry of Peptide Antibiotics*, H. Kleinkauf and H. von Dohren, editors, W. de Gruyter, Berlin, pp. 179-203), and also exhibit entomopathogenic properties (Grove and Pople, 1980, *Mycopathologia* 70: 103-105).

The enniatin synthetase gene (*esyn1*) has been cloned from *Fusarium scirpi* (Haese *et al.*, 1993, *Molecular Microbiology* 7: 905-914).

Enniatin synthetase mutants of *Fusarium avenaceum* have been generated that do not produce enniatins (Herrmann *et al.*, 1996, *Molecular Plant-Microbe Interactions* 9: 226-232).

It is an object of the present invention to provide methods for producing heterologous polypeptides in cyclohexadepsipeptide-deficient filamentous fungal mutant cells.

Summary of the Invention

The present invention relates to methods for producing a heterologous polypeptide, comprising: (a) cultivating a mutant of a parent filamentous fungal cell under conditions conducive for the production of the heterologous polypeptide, wherein (i) the mutant cell comprises a first nucleic acid sequence encoding the heterologous polypeptide, and (ii) the mutant produces less of a cyclohexadepsipeptide than the parent filamentous fungal cell when cultured under the same conditions; and (b) isolating the heterologous polypeptide from the cultivation medium.

The present invention also relates to mutants of filamentous fungal cells and methods

for obtaining the mutant cells.

The present invention also relates to isolated cyclohexadepsipeptide synthetases from *Fusarium venenatum* and isolated nucleic acid sequences encoding the cyclohexadepsipeptide synthetases. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the nucleic acid sequences as well as methods for producing the cyclohexadepsipeptide synthetases.

The present invention further relates to cyclohexadepsipeptides produced by the cyclohexadepsipeptide synthetases.

Brief Description of Figure

Figure 1 shows the genomic nucleic acid sequence and the deduced amino acid sequence of a *Fusarium venenatum* ATCC 20334 cyclohexadepsipeptide synthetase (SEQ ID NOS: 1 and 2, respectively).

Figure 2 shows the construction of pAES-*amdS*.

Detailed Description of the Invention

The present invention relates to methods for producing a heterologous polypeptide, comprising: (a) cultivating a mutant of a parent filamentous fungal cell under conditions conducive for the production of the heterologous polypeptide, wherein (i) the mutant filamentous fungal cell comprises a first nucleic acid sequence encoding the heterologous polypeptide and (ii) the mutant produces less of a cyclohexadepsipeptide than the parent filamentous fungal cell when cultured under the same conditions; and (b) isolating the heterologous polypeptide from the cultivation medium of the mutant cell.

The term "cyclohexadepsipeptide" is defined herein as a family of peptide-related compounds composed of hydroxy and amino acids linked by amide and ester bonds.

The term "production of a cyclohexadepsipeptide" is defined herein as to include any step involved in the production of a cyclohexadepsipeptide including, but not limited to, biosynthesis, regulation of biosynthesis, transport, and secretion.

In a preferred embodiment, the cyclohexadepsipeptide is an enniatin.

The term "enniatins" is defined herein as a family of cyclohexadepsipeptides

composed of three D-2-hydroxyisovaleric acid residues joined alternatively to L-amino acids or N-methyl-L-amino acids to produce an 18-membered cyclic structure. The ennatiins include, but are not limited to, ennatin A, A₁, B, B₁, B₂, B₃, B₄, C, D, E, and F; and derivatives thereof (Visconte *et al.*, 1992, *Journal of Agricultural and Food Chemistry* 40: 1076-1082; Tomodo *et al.*, 1992, *Journal of Antibiotics* 45: 1207-1215), and mixed-type ennatiins containing more than one species of amino acid (Zocher *et al.* 1982, *Biochemistry* 21: 43-48).

In the methods of the present invention, the filamentous fungal cell may be a wild-type cell or a mutant thereof. Furthermore, the filamentous fungal cell may be a cell that does not produce any detectable cyclohexadepsipeptide(s), but contains the genes encoding the cyclohexadepsipeptide(s). Preferably, the filamentous fungal cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Beauveria*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Gibberella*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Myrothecium*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Piromyces*, *Polyporus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, or *Trichoderma* cell.

In a preferred embodiment, the filamentous fungal cell is an *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, or *Aspergillus oryzae* cell.

In another preferred embodiment, the filamentous fungal cell is a *Fusarium acuminatum*, *Fusarium avenaceum*, *Fusarium bacridioides*, *Fusarium compactum*, *Fusarium crookwellense* (synonym of *Fusarium cerealis*), *Fusarium culmorum*, *Fusarium equiseti*, *Fusarium gibbosum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium lateritium*, *Fusarium moniliforme*, *Fusarium negundi*, *Fusarium nivale*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcocrochrum*, *Fusarium scirpi*, *Fusarium semitectum*, *Fusarium solani*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium tricinctum*, or *Fusarium venenatum* cell.

In another preferred embodiment, the filamentous fungal cell is a *Gibberella pulicaris*, *Gibberella zeae*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Myrothecium roridin*, *Neurospora crassa*, *Paecilomyces fumoso-roseus*, *Penicillium purpurogenum*, or *Polyporus sulphureus* cell.

In another preferred embodiment, the filamentous fungal cell is a *Trichoderma*

harzianum, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

In a more preferred embodiment, the *Fusarium venenatum* cell is *Fusarium venenatum* A3/5, which was originally deposited as *Fusarium graminearum* ATCC 20334 and recently reclassified as *Fusarium venenatum* by Yoder and Christianson, 1998, *Fungal Genetics and Biology* 23: 62-80 and O'Donnell et al., 1998, *Fungal Genetics and Biology* 23: 57-67; as well as taxonomic equivalents of *Fusarium venenatum* regardless of the species name by which they are currently known. In another more preferred embodiment, the *Fusarium venenatum* cell is a morphological mutant of *Fusarium venenatum* A3/5 or *Fusarium venenatum* ATCC 20334, as disclosed in WO 97/26330.

The filamentous fungal cell may also be a cell involved in the production of products containing (parts of) the mycelium, for example, in the production of the product QUORN™ (Marlow Foods, Ltd., Great Britain), which is produced from a *Fusarium* strain.

In the methods of the present invention, the mutant cell comprises a second nucleic acid sequence which comprises a modification of at least one of the genes involved in the production of the cyclohexadepsipeptide. Any gene of a filamentous fungal cell involved in the production of a cyclohexadepsipeptide may be modified. In a preferred embodiment, the gene is a cyclohexadepsipeptide synthetase gene. In a more preferred embodiment, the gene is an ennatin synthetase gene. In another more preferred embodiment, the gene is a D-hydroxyisovalerate dehydrogenase gene. D-Hydroxyisovalerate dehydrogenase catalyzes the conversion of 2-ketoisovalerate to D-hydroxyisovalerate (Lee and Zocher, 1996, *Journal of Biochemistry and Molecular Biology* 29: 493-499). In an even more preferred embodiment, the gene is a *Fusarium venenatum* cyclohexadepsipeptide synthetase gene having (a) a nucleic acid sequence encoding a polypeptide having an amino acid sequence which has at least 65% identity with the mature polypeptide contained within SEQ ID NO:2; (b) a nucleic acid sequence having at least 65% homology with the mature polypeptide coding region of SEQ ID NO:1; (c) a nucleic acid sequence which hybridizes under medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii) of at least 100 nucleotides, or (iv) a complementary strand of (i), (ii), or (iii); (d) an allelic variant of (a), (b), or (c); or (e) a subsequence of (a), (b), (c), or (d), wherein the subsequence encodes a polypeptide fragment which has cyclohexadepsipeptide synthetase activity. In a most preferred embodiment, the

gene is a *Fusarium venenatum* cyclohexadepsipeptide synthetase gene having the nucleic acid sequence of SEQ ID NO:1.

The cyclohexadepsipeptide-deficient filamentous fungal mutant cell may be constructed by reducing or eliminating expression of one or more of the genes described
5 above using methods well known in the art, for example, insertions, disruptions, replacements, or deletions. The gene to be modified or inactivated may be, for example, the coding region or a part thereof essential for activity, or a regulatory element of the gene required for the expression of the coding region. An example of such a regulatory or control
10 sequence may be a promoter sequence or a functional part thereof, *i.e.*, a part that is sufficient for affecting expression of the nucleic acid sequence. Other control sequences for possible modification include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, signal peptide sequence, transcription terminator, and transcriptional activator.

Modification or inactivation of the gene may be performed by subjecting the parent cell to mutagenesis and selecting for mutant cells in which expression of the gene has been
15 reduced or eliminated. The mutagenesis, which may be specific or random, may be performed, for example, by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the mutagenesis may be performed by use of any combination of these mutagenizing agents.

Examples of a physical or chemical mutagenizing agent suitable for the present
20 purpose include ultraviolet (UV) irradiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues.

When such agents are used, the mutagenesis is typically performed by incubating the
25 parent cell to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions, and screening and/or selecting for mutant cells exhibiting reduced or no expression of the gene.

Modification or inactivation of the gene may be accomplished by introduction, substitution, or removal of one or more nucleotides in the gene or a regulatory element
30 required for the transcription or translation thereof. For example, nucleotides may be inserted or removed so as to result in the introduction of a stop codon, the removal of the start codon, or a change in the open reading frame. Such modification or inactivation may be

accomplished by site-directed mutagenesis or PCR generated mutagenesis in accordance with methods known in the art. Although, in principle, the modification may be performed *in vivo*, *i.e.*, directly on the cell expressing the gene to be modified, it is preferred that the modification be performed *in vitro* as exemplified below.

5 An example of a convenient way to eliminate or reduce production of a cyclohexadepsipeptide by a filamentous fungal cell of choice is based on techniques of gene replacement, gene deletion, or gene disruption. For example, in the gene disruption method, a nucleic acid sequence corresponding to the endogenous gene or gene fragment of interest is mutagenized *in vitro* to produce a defective nucleic acid sequence which is then transformed
10 into the parent cell to produce a defective gene. By homologous recombination, the defective nucleic acid sequence replaces the endogenous gene or gene fragment. It may be desirable that the defective gene or gene fragment also encodes a marker that may be used for selection of transformants in which the nucleic acid sequence has been modified or destroyed. In a particularly preferred embodiment, the gene is disrupted with a selectable marker such as those described herein.

15 Alternatively, modification or inactivation of the gene may be performed by established anti-sense techniques using a nucleotide sequence complementary to the nucleic acid sequence of the gene. More specifically, expression of the gene by a filamentous fungal cell may be reduced or eliminated by introducing a nucleotide sequence complementary to the nucleic acid sequence of the gene that may be transcribed in the cell and is capable of
20 hybridizing to the mRNA produced in the cell. Under conditions allowing the complementary anti-sense nucleotide sequence to hybridize to the mRNA, the amount of protein translated is thus reduced or eliminated.

25 A nucleic acid sequence complementary or homologous to the nucleic acid sequence of a gene involved in the production of a cyclohexadepsipeptide may be obtained from other microbial sources that produce cyclohexadepsipeptides.

Preferred sources for an enniatin synthetase gene having a nucleic acid sequence complementary or homologous to the nucleic acid sequence of SEQ ID NO:1 of *Fusarium venenatum* include other *Fusarium* strains. A more preferred source is *Fusarium scirpi*
30 (Haese *et al.*, 1993, *supra*).

Preferred sources for D-hydroxyisovalerate dehydrogenase genes that may be complementary or homologous to the nucleic acid sequence of the corresponding genes of a

filamentous fungal cell include other *Fusarium* strains. A more preferred source for the D-hydroxyisovalerate dehydrogenase gene is *Fusarium sambucinum* (Lee and Zocher, 1996, *supra*). Furthermore, the nucleic acid sequences may be native to the filamentous fungal cell.

The level of cyclohexadepsipeptides produced by a mutant filamentous fungal cell of the present invention may be determined using the method of Visconti *et al.*, 1992, *Journal of Agriculture and Food Chemistry* 40: 1076-1082. Specifically, one ml of *Fusarium venenatum* cell-free culture broth is extracted twice with 2.0 ml ethyl acetate. The combined organic extracts are evaporated to dryness under a stream of nitrogen gas and redissolved in 0.5 ml hexane. One microliter samples are analyzed using a Hewlett-Packard 6890 GC/MSD system operating in the electron impact (EI) mode. Samples are injected on-column and separated utilizing a DB-5 capillary column (30 m x 0.25 mm, 0.25 μ m film) employing a temperature program with heating from 120 to 300°C at a rate of 15°C/min. For example, enniatins A, A1, B, B1, B2 and B3 are identified by *m/z* ratios for the (M^+ + H) ion of 682, 668, 640, 654, 626 and 612, respectively.

The mutant filamentous fungal cell preferably produces at least about 25% less, more preferably at least about 50% less, even more preferably at least about 75% less, most preferably at least about 95% less, and even most preferably no cyclohexadepsipeptide than the corresponding parent filamentous fungal cell when cultured under identical conditions. The parent and mutant cells may be compared with regard to production of a cyclohexadepsipeptide under conditions conducive for the production of a polypeptide of interest or under conditions conducive for the production of a cyclohexadepsipeptide.

In another aspect of the present invention, the mutant filamentous fungal cell may additionally contain modifications of one or more third nucleic acid sequences that encode proteins that may be detrimental to the production, recovery, and/or application of the heterologous polypeptide of interest. The modification reduces or eliminates expression of the one or more third nucleic acid sequences resulting in a mutant cell that may produce more of the heterologous polypeptide than the mutant cell without the modification of the third nucleic acid sequence when cultured under the same conditions. The third nucleic acid sequence may encode any protein or enzyme. For example, the enzyme may be an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, invertase, laccase,

lipase, mannosidase, mutanase, oxidase, pectinolytic enzyme, peroxidase, phospholipase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or xylanase. The third nucleic acid sequence preferably encodes a proteolytic enzyme, e.g., an aminopeptidase, carboxypeptidase, or endoprotease.

5 The mutant filamentous fungal cell is cultivated in a nutrient medium suitable for production of a heterologous polypeptide of interest using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions
10 allowing the heterologous polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). The secreted heterologous polypeptide can be recovered
15 directly from the medium.

The heterologous polypeptide may be detected using methods known in the art that are specific for the polypeptide. These detection methods may include use of specific antibodies, formation of an enzyme product, disappearance of an enzyme substrate, or SDS-PAGE. For example, an enzyme assay may be used to determine the activity of the heterologous polypeptide. Procedures for determining enzyme activity are known in the art
20 for many enzymes.

The resulting heterologous polypeptide may be isolated by methods known in the art. For example, the polypeptide may be isolated from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray-drying,
25 evaporation, or precipitation. The isolated polypeptide may then be further purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), or extraction (see, e.g., *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989).
30

The polypeptide may be any polypeptide heterologous to the mutant filamentous fungal cell. The term "polypeptide" is not meant herein to refer to a specific length of the

002110 00220160
5 encoded product and, therefore, encompasses peptides, oligopeptides, and proteins. The term "heterologous polypeptide" is defined herein as a polypeptide that is not native to the fungal cell, a native protein in which modifications have been made to alter the native sequence, or a native protein whose expression is quantitatively altered as a result of a manipulation of the fungal cell by recombinant DNA techniques. The mutant fungal cell may contain one or more copies of the nucleic acid sequence encoding the polypeptide. In a preferred embodiment, the heterologous polypeptide is an extracellularly secreted polypeptide.

10 Preferably, the heterologous polypeptide is a hormone, hormone variant, enzyme, receptor or portion thereof, antibody or portion thereof, or reporter. In a more preferred embodiment, the heterologous polypeptide is an oxidoreductase, transferase, hydrolase, lyase, isomerase, or ligase. In an even more preferred embodiment, the heterologous polypeptide is an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, invertase, laccase, lipase, mannosidase, mutanase, oxidase, pectinolytic enzyme, peroxidase, phospholipase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase or xylanase.

15 The nucleic acid sequence encoding a heterologous polypeptide that can be expressed in a filamentous fungal cell may be obtained from any prokaryotic, eukaryotic, or other source. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide is produced by the source or by a cell in which a gene from the source has been inserted.

20 In the methods of the present invention, the mutant filamentous fungal cell may also be used for the recombinant production of polypeptides that are native to the cell. The native polypeptides may be recombinantly produced by, e.g., placing a gene encoding the polypeptide under the control of a different promoter to enhance expression of the polypeptide, to expedite export of a native polypeptide of interest outside the cell by use of a signal sequence, and to increase the copy number of a gene encoding the polypeptide normally produced by the cell. The present invention also encompasses, within the scope of the term "heterologous polypeptide", such recombinant production of homologous polypeptides, to the extent that such expression involves the use of genetic elements not native to the cell, or use of native elements that have been manipulated to function in a manner that do not normally occur in the host cell.

The techniques used to isolate or clone a nucleic acid sequence encoding a heterologous polypeptide are known in the art and include isolation from genomic DNA, preparation from cDNA, or a combination thereof. The cloning of the nucleic acid sequence from such genomic DNA can be effected, e.g., by using the well known polymerase chain
5 reaction (PCR). See, for example, Innis *et al.*, 1990, *PCR Protocols: A Guide to Methods and Application*, Academic Press, New York. The cloning procedures may involve excision and isolation of a desired nucleic acid fragment comprising the nucleic acid sequence encoding the polypeptide, insertion of the fragment into a vector molecule, and incorporation of the recombinant vector into the mutant fungal cell where multiple copies or clones of the
10 nucleic acid sequence will be replicated. The nucleic acid sequence may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

In the methods of the present invention, heterologous polypeptides may also include fused or hybrid polypeptides in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide or fragment thereof. A fused polypeptide is produced by
15 fusing a nucleic acid sequence (or a portion thereof) encoding one polypeptide to a nucleic acid sequence (or a portion thereof) encoding another polypeptide. Techniques for producing fusion polypeptides are known in the art, and include, ligating the coding sequences encoding the polypeptides so that they are in frame and expression of the fused polypeptide is under control of the same promoter(s) and terminator. The hybrid polypeptides may comprise a
20 combination of partial or complete polypeptide sequences obtained from at least two different polypeptides wherein one or more may be heterologous to the mutant filamentous fungal cell.

An isolated nucleic acid sequence encoding a heterologous polypeptide of interest may be manipulated in a variety of ways to provide for expression of the polypeptide. Expression will be understood to include any step involved in the production of the
25 polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion. Manipulation of the nucleic acid sequence prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying nucleic acid sequences utilizing cloning methods are well known in the art.

"Nucleic acid construct" is defined herein as a nucleic acid molecule, either single- or
30 double-stranded, isolated from a naturally occurring gene or modified to contain segments of nucleic acid that are combined and juxtaposed in a manner which would not otherwise exist

in nature. The term nucleic acid construct is synonymous with the term expression cassette when the nucleic acid construct contains all the control sequences required for expression of a coding sequence. The term "coding sequence" as defined herein is a sequence that is transcribed into mRNA and translated into a polypeptide. The boundaries of the coding
5 sequence are generally determined by the ATG start codon located just upstream of the open reading frame at the 5' end of the mRNA and a transcription terminator sequence located just downstream of the open reading frame at the 3' end of the mRNA. A coding sequence can include, but is not limited to, genomic, cDNA, RNA, semisynthetic, synthetic, recombinant, or any combinations thereof.

10 The term "control sequences" is defined herein to include all components that are necessary or advantageous for the expression of a heterologous polypeptide. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a
15 minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleic acid sequence encoding a heterologous polypeptide. The term "operably linked" is defined herein as a configuration in which a control sequence is appropriately placed at a
20 position relative to the coding sequence of the DNA sequence such that the control sequence directs the production of a heterologous polypeptide.

The control sequence may be an appropriate promoter sequence, a nucleic acid sequence that is recognized by a filamentous fungal cell for expression of the nucleic acid sequence. The promoter sequence contains transcriptional control sequences that mediate the
25 expression of the heterologous polypeptide. The promoter may be any nucleic acid sequence that shows transcriptional activity in the filamentous fungal cell including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the cell.

30 Examples of suitable promoters for directing the transcription of the nucleic acid constructs in the methods of the present invention are promoters obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase,

004482788-01300

Aspergillus niger or *Aspergillus awamori* glucoamylase (*glaA*), *Rhizomucor miehei* lipase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Aspergillus nidulans* acetamidase, *Aspergillus oryzae* acetamidase (*amdS*), *Fusarium oxysporum* trypsin-like protease (U.S. Patent No. 4,288,627), and mutant, truncated, and
5 hybrid promoters thereof. Particularly preferred promoters are the NA2-tpi (a hybrid of the promoters from the genes encoding *Aspergillus niger* neutral alpha-amylase and *Aspergillus oryzae* triose phosphate isomerase), glucoamylase, and TAKA amylase promoters.

The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a filamentous fungal cell to terminate transcription. The terminator
10 sequence is operably linked to the 3' terminus of the nucleic acid sequence encoding the heterologous polypeptide. Any terminator that is functional in the filamentous fungal cell may be used in the present invention.

Preferred terminators are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthetase, *Aspergillus niger* alpha-glucosidase, and *Fusarium oxysporum* trypsin-like protease.
15

The control sequence may also be a suitable leader sequence, a nontranslated region of a mRNA that is important for translation by the filamentous fungal cell. The leader sequence is operably linked to the 5' terminus of the nucleic acid sequence encoding the heterologous polypeptide. Any leader sequence that is functional in the filamentous fungal
20 cell may be used in the present invention.

Preferred leaders are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3' terminus of the nucleic acid sequence and, when transcribed, is recognized by
25 a filamentous fungal cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the filamentous fungal cell may be used in the present invention.

Preferred polyadenylation sequences are obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* glucoamylase, *Aspergillus nidulans* anthranilate synthase, and *Aspergillus niger* alpha-glucosidase.
30

The control sequence may also be a signal peptide coding region that codes for an amino acid sequence linked to the amino terminus of the heterologous polypeptide and directs

the encoded polypeptide into the cell's secretory pathway. The 5' end of the coding sequence of the nucleic acid sequence may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region that encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding region that is foreign to the coding sequence. The foreign signal peptide coding region may be required where the coding sequence does not naturally contain a signal peptide coding region. Alternatively, the foreign signal peptide coding region may simply replace the natural signal peptide coding region in order to enhance secretion of the polypeptide. The signal peptide coding region may be obtained from a glucoamylase or an amylase gene from an *Aspergillus* species, or a lipase or proteinase gene from a *Rhizomucor* species. However, any signal peptide coding region that directs the expressed heterologous polypeptide into the secretory pathway of a filamentous fungal cell may be used in the present invention.

An effective signal peptide coding region is the signal peptide coding region obtained from the genes encoding *Aspergillus oryzae* TAKA amylase, *Aspergillus niger* neutral amylase, *Rhizomucor miehei* aspartic proteinase gene, and *Humicola lanuginosa* cellulase.

The control sequence may also be a propeptide coding region that codes for an amino acid sequence positioned at the amino terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to a mature, active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding region may be obtained from the genes encoding *Rhizomucor miehei* aspartic proteinase and *Myceliophthora thermophila* laccase (WO 95/33836).

Where both signal peptide and propeptide regions are present at the amino terminus of a polypeptide, the propeptide region is positioned next to the amino terminus of the polypeptide and the signal peptide region is positioned next to the amino terminus of the propeptide region.

The nucleic acid constructs may also comprise one or more nucleic acid sequences that encode one or more factors that are advantageous for directing the expression of the heterologous polypeptide, e.g., a transcriptional activator (e.g., a *trans*-acting factor), chaperone, and processing protease. Any factor that is functional in a filamentous fungal cell may be used in the present invention. The nucleic acids encoding one or more of these

003710-88228460

factors are not necessarily in tandem with the nucleic acid sequence encoding the heterologous polypeptide.

It may also be desirable to add regulatory sequences that allow the regulation of the expression of the heterologous polypeptide relative to the growth of the filamentous fungal cell. Examples of regulatory systems are those that cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. The TAKA alpha-amylase promoter, *Aspergillus niger* glucoamylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used as regulatory sequences. Other examples of regulatory sequences are those that allow for gene amplification, e.g., the metallothionein genes which are amplified with heavy metals. In these cases, the nucleic acid sequence encoding the heterologous polypeptide would be operably linked with the regulatory sequence.

The various nucleic acid and control sequences described above may be joined together to produce a recombinant expression vector that may include one or more convenient restriction sites to allow for insertion or substitution of the nucleic acid sequence encoding the heterologous polypeptide at such sites. Alternatively, the nucleic acid sequence encoding the heterologous polypeptide may be expressed by inserting the sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression, and possibly secretion.

The recombinant expression vector may be any vector (e.g., a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about the expression of the nucleic acid sequence encoding the heterologous polypeptide. The choice of the vector will typically depend on the compatibility of the vector with the filamentous fungal cell into which the vector is to be introduced. The vector may be a linear or closed circular plasmid. The vector may be an autonomously replicating vector, i.e., a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the filamentous fungal cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. The vector system may be a single vector or plasmid or two or more vectors or

plasmids that together contain the total DNA to be introduced into the genome of the filamentous fungal cell, or a transposon.

The vector preferably contains one or more selectable markers that permit easy selection of transformed filamentous fungal cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like. A selectable marker for use in a filamentous fungal host cell may be selected from the group including, but not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hygB* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in a filamentous fungal cell are the *amdS* and *pyrG* genes of *Aspergillus nidulans* or *Aspergillus oryzae* and the *bar* gene of *Streptomyces hygroscopicus*.

The vector preferably contains an element(s) that permits stable integration of the vector into a filamentous fungal cell genome or autonomous replication of the vector in the cell independent of the genome of the cell.

"Introduction" means introducing a vector comprising the nucleic acid sequence into a filamentous fungal cell so that the vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector. Integration is generally considered to be an advantage as the nucleic acid sequence is more likely to be stably maintained in the cell. Integration of the vector into the chromosome occurs by homologous recombination, non-homologous recombination, or transposition.

The introduction of an expression vector into a filamentous fungal cell may involve a process consisting of protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* host cells are described in EP 238 023 and Yelton *et al.*, 1984, *Proceedings of the National Academy of Sciences USA* 81: 1470-1474. A suitable method of transforming *Fusarium* species is described by Malardier *et al.*, 1989, *Gene* 78: 147-156 or in WO 96/00787.

For integration into the genome of a filamentous fungal cell, the vector may rely on the nucleic acid sequence encoding the heterologous polypeptide or any other element of the vector for stable integration of the vector into the genome by homologous or nonhomologous recombination. Alternatively, the vector may contain additional nucleic acid sequences for

directing integration by homologous recombination into the genome of the filamentous fungal cell. The additional nucleic acid sequences enable the vector to be integrated into the genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should preferably contain a sufficient number of nucleic acids, such as 100 to 1,500 base pairs, preferably 400 to 1,500 base pairs, and most preferably 800 to 1,500 base pairs, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequences that are homologous with the target sequence in the genome of the filamentous fungal cell. Furthermore, the integrational elements may be non-encoding or encoding nucleic acid sequences. On the other hand, the vector may be integrated into the genome of the cell by non-homologous recombination.

For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the filamentous fungal cell in question.

It will be understood that the methods of the present invention are not limited to a particular order for obtaining the mutant filamentous fungal cell. The modification of a gene involved in the production of a cyclohexadepsipeptide may be introduced into the parent cell at any step in the construction of the cell for the production of a heterologous polypeptide. It is preferable that the filamentous fungal mutant has already been made cyclohexadepsipeptide-deficient using the methods of the present invention prior to the introduction of a gene encoding a heterologous polypeptide.

The procedures used to ligate the elements described herein to construct the recombinant expression vectors are well known to one skilled in the art (see, e.g., J. Sambrook, E.F. Fritsch, and T. Maniatus, 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York).

The present invention also relates to methods for obtaining cyclohexadepsipeptide-deficient filamentous fungal mutant cells which comprise (a) introducing into a parent filamentous fungal cell a first nucleic acid sequence comprising a modification of at least one of the genes involved in the production of a cyclohexadepsipeptide and a second nucleic acid sequence encoding a heterologous polypeptide; and (b) identifying the mutant from step (a) comprising the modified nucleic acid sequence, wherein the mutant cell produces less of the cyclohexadepsipeptide than the parent filamentous fungal cell of the mutant cell when cultured under the same conditions.

The present invention also relates to cyclohexadepsipeptide-deficient mutants of filamentous fungal cells for producing a heterologous polypeptide which comprise a first nucleic acid sequence comprising a modification of at least one of the genes involved in the production of a cyclohexadepsipeptide and a second nucleic acid sequence encoding the heterologous polypeptide, wherein the mutant produces less of the cyclohexadepsipeptide than the parent filamentous fungal cell of the mutant cell when cultured under the same conditions.

The present invention also relates to isolated cyclohexadepsipeptide synthetases. The term "cyclohexadepsipeptide synthetase activity" is defined herein as a synthetase activity which catalyzes the production of a cyclohexadepsipeptide from D-2-hydroxyisovaleric acid, a branched chain L-amino acid (e.g., valine, leucine, isoleucine), S-adenosylmethionine, and ATP. For purposes of the present invention, cyclohexadepsipeptide synthetase activity is determined by measuring the production of a cyclohexadepsipeptide according to the procedure of Zocher *et al.*, 1982, *Biochemistry* 21: 43-48. Specifically, the cyclohexadepsipeptide synthetase is incubated with 1 mM valine, 0.2 mM S-adenosylmethionine, 0.2 mM D-2-hydroxyisovaleric acid, 4 mM ATP, and 4 mM Mg(OAc)₂ in a total volume of 100 µl for 10 minutes at 37°C in 50 mM MOPS pH 7.0. The amount of cyclohexadepsipeptide is determined as described herein based on the method of Visconti *et al.*, 1992, *supra*. One unit of cyclohexadepsipeptide synthetase activity is defined as 1.0 µmole of cyclohexadepsipeptide produced per minute at 37°C, pH 7.0.

In a first embodiment, the present invention relates to isolated cyclohexadepsipeptide synthetases having an amino acid sequence which has a degree of identity to the mature polypeptide contained within SEQ ID NO:2 of at least about 65%, preferably at least about 70%, more preferably at least about 80%, even more preferably at least about 90%, most preferably at least about 95%, and even most preferably at least about 97%, which have cyclohexadepsipeptide synthetase activity (hereinafter "homologous cyclohexadepsipeptide synthetases"). In a preferred embodiment, the homologous cyclohexadepsipeptide synthetases have an amino acid sequence which differs by five amino acids, preferably by four amino acids, more preferably by three amino acids, even more preferably by two amino acids, and most preferably by one amino acid from the mature polypeptide contained within SEQ ID NO:2. For purposes of the present invention, the degree of identity between two amino acid sequences is determined by the Clustal method (Higgins, 1989, *CABIOS* 5: 151-

30

153) using the LASERGENE™ MEGALIGN™ software (DNASTAR, Inc., Madison, WI) with an identity table and the following multiple alignment parameters: Gap penalty of 10, and gap length penalty of 10. Pairwise alignment parameters were Ktuple=1, gap penalty=3, windows=5, and diagonals=5.

Preferably, the cyclohexadepsipeptide synthetases of the present invention comprise the amino acid sequence of SEQ ID NO:2 or an allelic variant thereof; or a fragment thereof that has cyclohexadepsipeptide synthetase activity. In a more preferred embodiment, the cyclohexadepsipeptide synthetase of the present invention comprises the amino acid sequence of SEQ ID NO:2. In another preferred embodiment, the cyclohexadepsipeptide synthetase of the present invention comprises the mature polypeptide contained within SEQ ID NO:2, or an allelic variant thereof; or a fragment thereof that has cyclohexadepsipeptide synthetase activity. In another preferred embodiment, the cyclohexadepsipeptide synthetase of the present invention comprises the mature polypeptide contained within SEQ ID NO:2. In another preferred embodiment, the cyclohexadepsipeptide synthetase of the present invention consists of the amino acid sequence of SEQ ID NO:2 or an allelic variant thereof; or a fragment thereof that has cyclohexadepsipeptide synthetase activity. In another preferred embodiment, the cyclohexadepsipeptide synthetase of the present invention consists of the amino acid sequence of SEQ ID NO:2. In another preferred embodiment, the cyclohexadepsipeptide synthetase consists of the mature polypeptide contained within SEQ ID NO:2 or an allelic variant thereof; or a fragment thereof that has cyclohexadepsipeptide synthetase activity. In another preferred embodiment, the cyclohexadepsipeptide synthetase consists of the mature polypeptide contained within SEQ ID NO:2.

A fragment of SEQ ID NO:2 is a polypeptide having one or more amino acids deleted from the amino and/or carboxyl terminus of this amino acid sequence. Preferably, a fragment contains at least 2854 amino acid residues, more preferably at least 2954 amino acid residues, and most preferably at least 3054 amino acid residues.

An allelic variant denotes any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

The amino acid sequences of the homologous cyclohexadepsipeptide synthetases may

00210-8822460

differ from the amino acid sequence of SEQ ID NO:2 or the mature polypeptide thereof by an insertion or deletion of one or more amino acid residues and/or the substitution of one or more amino acid residues by different amino acid residues. Preferably, amino acid changes are of a minor nature, that is conservative amino acid substitutions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

Examples of conservative substitutions are within the group of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter the specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, *In, The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly as well as these in reverse.

In a second embodiment, the present invention relates to isolated cyclohexadepsipeptide synthetases encoded by nucleic acid sequences that hybridize under low stringency conditions, more preferably medium stringency conditions, even more preferably high stringency conditions, and most preferably very high stringency conditions with a nucleic acid probe that hybridizes under the same conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii), or (iv) a complementary strand of (i), (ii), or (iii) (J. Sambrook, E.F. Fritsch, and T. Maniatus, 1989, *Molecular Cloning, A Laboratory Manual*, 2d edition, Cold Spring Harbor, New York). The subsequence of SEQ ID NO:1 may be at least 100 nucleotides or preferably at least 200 nucleotides. Moreover, the subsequence may encode a polypeptide fragment that has cyclohexadepsipeptide synthetase activity. The cyclohexadepsipeptide synthetases may also be allelic variants or fragments that have cyclohexadepsipeptide synthetase activity.

The nucleic acid sequence of SEQ ID NO:1 or a subsequence thereof, as well as the

amino acid sequence of SEQ ID NO:2 or a fragment thereof, may be used to design a nucleic acid probe to identify and clone DNA encoding cyclohexadepsipeptide synthetases from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic or cDNA of the genus or species of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 15, preferably at least 25, and more preferably at least 35 nucleotides in length. Longer probes can also be used. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ^{32}P , ^3H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present invention.

Thus, a genomic DNA or cDNA library prepared from such other organisms may be screened for DNA that hybridizes with the probes described above and encodes a cyclohexadepsipeptide synthetase. Genomic or other DNA from such other organisms may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques.

DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that is homologous with SEQ ID NO:1 or a subsequence thereof, the carrier material is used in a Southern blot. For purposes of the present invention, hybridization indicates that the nucleic acid sequence hybridizes to a nucleic acid probe corresponding to (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii), or (iv) a complementary strand of (i), (ii), or (iii) under low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under these conditions are detected using X-ray film.

In a preferred embodiment, the nucleic acid probe is a nucleic acid sequence that encodes the cyclohexadepsipeptide synthetase of SEQ ID NO:2, or a subsequence thereof. In another preferred embodiment, the nucleic acid probe is SEQ ID NO:1. In another preferred embodiment, the nucleic acid probe is the mature polypeptide coding region contained within SEQ ID NO:1. In another preferred embodiment, the nucleic acid probe is the nucleic acid sequences contained in plasmid pZL-ESA, which is contained in *Escherichia coli* NRRL B-30068, plasmid pZL-ESB, which is contained in *Escherichia coli* NRRL B-30069, and plasmid pZL-ESC, which is contained in *Escherichia coli* NRRL B-30070, wherein the nucleic acid sequences encode the cyclohexadepsipeptide synthetase of SEQ ID NO:2. In

another preferred embodiment, the nucleic acid probe is the nucleic acid sequence encoding the mature cyclohexadepsipeptide synthetase of SEQ ID NO:2 contained in plasmid pZL-ESA, which is contained in *Escherichia coli* NRRL B-30068, plasmid pZL-ESB, which is contained in *Escherichia coli* NRRL B-30069, and plasmid pZL-ESC, which is contained in
5 *Escherichia coli* NRRL B-30070.

For long probes of at least 100 nucleotides in length, low to very high stringency conditions are defined as prehybridization and hybridization at 45°C in 5X SSPE, 0.3% SDS, 200 µg/ml sheared and denatured salmon sperm DNA, and either 25% formamide for very low and low stringencies, 35% formamide for medium and medium-high stringencies, or 50%
10 formamide for high and very high stringencies, following standard Southern blotting procedures.

For long probes of at least 100 nucleotides in length, the carrier material is finally washed three times each for 15 minutes using 2 x SSC, 0.2% SDS preferably at least at 45°C (very low stringency), more preferably at least at 50°C (low stringency), more preferably at least at 55°C (medium stringency), more preferably at least at 60°C (medium-high stringency), even more preferably at least at 65°C (high stringency), and most preferably at least at 70°C (very high stringency).

For short probes that are about 15 nucleotides to about 70 nucleotides in length, stringency conditions are defined as prehybridization, hybridization, and washing post-hybridization at 5°C to 10°C below the calculated T_m using the calculation according to Bolton and McCarthy (1962, *Proceedings of the National Academy of Sciences USA* 48:1390) in 0.9 M NaCl, 0.09 M Tris-HCl pH 7.6, 6 mM EDTA, 0.5% NP-40, 1X Denhardt's solution, 1 mM sodium pyrophosphate, 1 mM sodium monobasic phosphate, 0.1 mM ATP, and 0.2 mg
20 of yeast RNA per ml following standard Southern blotting procedures.

For short probes that are about 15 nucleotides to about 70 nucleotides in length, the carrier material is washed once in 6X SSC plus 0.1% SDS for 15 minutes and twice each for 15 minutes using 6X SSC at 5°C to 10°C below the calculated T_m .
25

In a third embodiment, the present invention relates to isolated polypeptides having immunochemical identity or partial immunochemical identity to the cyclohexadepsipeptide
30 synthetase having the amino acid sequence of SEQ ID NO:2 or the mature polypeptide thereof. The immunochemical properties are determined by immunological cross-reaction identity tests by the well-known Ouchterlony double immunodiffusion procedure.

Specifically, an antiserum containing polyclonal antibodies that are immunoreactive or bind to epitopes of the polypeptide having the amino acid sequence of SEQ ID NO:2 or the mature polypeptide thereof are prepared by immunizing rabbits (or other rodents) according to the procedure described by Harboe and Ingild, *In N.H. Axelsen, J. Krøll, and B. Weeks, editors, A Manual of Quantitative Immunelectrophoresis*, Blackwell Scientific Publications, 1973, Chapter 23, or Johnstone and Thorpe, *Immunochemistry in Practice*, Blackwell Scientific Publications, 1982 (more specifically pages 27-31). A polypeptide having immunochemical identity is a polypeptide that reacts with the antiserum in an identical fashion such as total fusion of precipitates, identical precipitate morphology, and/or identical electrophoretic mobility using a specific immunochemical technique. A further explanation of immunochemical identity is described by Axelsen, Bock, and Krøll, *In N.H. Axelsen, J. Krøll, and B. Weeks, editors, A Manual of Quantitative Immunelectrophoresis*, Blackwell Scientific Publications, 1973, Chapter 10. A polypeptide having partial immunochemical identity is a polypeptide that reacts with the antiserum in a partially identical fashion such as partial fusion of precipitates, partially identical precipitate morphology, and/or partially identical electrophoretic mobility using a specific immunochemical technique. A further explanation of partial immunochemical identity is described by Bock and Axelsen, *In N.H. Axelsen, J. Krøll, and B. Weeks, editors, A Manual of Quantitative Immunelectrophoresis*, Blackwell Scientific Publications, 1973, Chapter 11.

The antibody may also be a monoclonal antibody. Monoclonal antibodies may be prepared and used, *e.g.*, according to the methods of E. Harlow and D. Lane, editors, 1988, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, New York.

The isolated cyclohexadepsipeptide synthetases of the present invention have at least 20%, preferably at least 40%, more preferably at least 60%, even more preferably at least 80%, even more preferably at least 90%, and most preferably at least 100% of the cyclohexadepsipeptide synthetase activity of the mature polypeptide of SEQ ID NO:2.

In a preferred embodiment, a cyclohexadepsipeptide synthetase of the present invention is obtained from a *Fusarium venenatum* strain, and more preferably from *Fusarium venenatum* ATCC 20334 or a mutant strain thereof, *e.g.*, the polypeptide with the amino acid sequence of SEQ ID NO:2.

As defined herein, an "isolated" cyclohexadepsipeptide synthetase is a polypeptide

that is essentially free of other polypeptides, e.g., at least about 20% pure, preferably at least about 40% pure, more preferably about 60% pure, even more preferably about 80% pure, most preferably about 90% pure, and even most preferably about 95% pure, as determined by SDS-PAGE.

5 A cyclohexadepsipeptide synthetase of the present invention may be obtained from microorganisms of any genus.

A cyclohexadepsipeptide synthetase of the present invention may be a bacterial cyclohexadepsipeptide synthetase. For example, the cyclohexadepsipeptide synthetase may be a gram positive bacterial cyclohexadepsipeptide synthetase such as a *Bacillus* cyclohexadepsipeptide synthetase, e.g., a *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*,
10 *Bacillus brevis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or *Bacillus thuringiensis* cyclohexadepsipeptide synthetase; or a *Streptomyces* cyclohexadepsipeptide synthetase, e.g., a *Streptomyces lividans* or *Streptomyces murinus* cyclohexadepsipeptide synthetase; or a gram negative bacterial cyclohexadepsipeptide synthetase, e.g., an *E. coli* or a *Pseudomonas* sp. cyclohexadepsipeptide synthetase.

A cyclohexadepsipeptide synthetase of the present invention may be a fungal cyclohexadepsipeptide synthetase, and more preferably a yeast cyclohexadepsipeptide synthetase such as a *Candida*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*,
15 or *Yarrowia* cyclohexadepsipeptide synthetase; or more preferably a filamentous fungal cyclohexadepsipeptide synthetase such as an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Piromyces*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, or *Trichoderma*
20 cyclohexadepsipeptide synthetase.

In a preferred embodiment, the cyclohexadepsipeptide synthetase is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis* or *Saccharomyces oviformis* cyclohexadepsipeptide synthetase.

30 In another preferred embodiment, the cyclohexadepsipeptide synthetase is an *Aspergillus aculeatus*, *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Humicola insolens*, *Humicola*

lanuginosa, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cyclohexadepsipeptide synthetase.

In another preferred embodiment, the cyclohexadepsipeptide synthetase is a *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, or *Fusarium venenatum* cyclohexadepsipeptide synthetase.

The present invention also relates to isolated nucleic acid sequences that encode a cyclohexadepsipeptide synthetase of the present invention. In a preferred embodiment, the nucleic acid sequence is set forth in SEQ ID NO:1. In another more preferred embodiment, the nucleic acid sequence is the sequences contained in plasmid pZL-ESA, which is contained in *Escherichia coli* NRRL B-30068, plasmid pZL-ESB, which is contained in *Escherichia coli* NRRL B-30069, and plasmid pZL-ESC, which is contained in *Escherichia coli* NRRL B-30070. In another more preferred embodiment, the nucleic acid sequence is the sequences encoding the mature polypeptide contained within SEQ ID NO:2 that is contained in plasmid pZL-ESA, which is contained in *Escherichia coli* NRRL B-30068, plasmid pZL-ESB, which is contained in *Escherichia coli* NRRL B-30069, and plasmid pZL-ESC, which is contained in *Escherichia coli* NRRL B-30070. In another preferred embodiment, the nucleic acid sequence is the mature polypeptide coding region contained within SEQ ID NO:1. The present invention also encompasses nucleic acid sequences that encode a polypeptide having the amino acid sequence of SEQ ID NO:2 or the mature polypeptide thereof, which differ from SEQ ID NO:1 by virtue of the degeneracy of the genetic code. The present invention also relates to subsequences of SEQ ID NO:1 that encode fragments of SEQ ID NO:2, which have cyclohexadepsipeptide synthetase activity.

A subsequence of SEQ ID NO:1 is a nucleic acid sequence encompassed by SEQ ID NO:1 except that one or more nucleotides from the 5' and/or 3' end have been deleted. Preferably, a subsequence contains at least 8562 nucleotides, more preferably at least 8862 nucleotides, and most preferably at least 9162 nucleotides.

The present invention also relates to mutant nucleic acid sequences comprising at least

one mutation in the mature polypeptide coding sequence of SEQ ID NO:1, in which the mutant nucleic acid sequence encodes a polypeptide which consists of the mature polypeptide contained within SEQ ID NO:2.

The techniques used to isolate or clone a nucleic acid sequence encoding a cyclohexadepsipeptide synthetase may include isolation from genomic DNA, preparation from cDNA, or a combination thereof, as described herein. The cloning of the nucleic acid sequences of the present invention from such genomic DNA can be effected, *e.g.*, by PCR or other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligated activated transcription (LAT) and nucleic acid sequence-based amplification (NASBA) may be used. The nucleic acid sequence may be cloned from a strain of *Fusarium*, or another or related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the nucleic acid sequence. The nucleic acid sequence may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

The term "isolated nucleic acid sequence" as used herein refers to a nucleic acid sequence that is essentially free of other nucleic acid sequences, *e.g.*, at least about 20% pure, preferably at least about 40% pure, more preferably at least about 60% pure, even more preferably at least about 80% pure, and most preferably at least about 90% pure as determined by agarose electrophoresis.

The present invention also relates to nucleic acid sequences that have a degree of homology to the mature polypeptide coding region contained within SEQ ID NO:1 of at least about 65%, preferably about 70%, preferably about 80%, more preferably about 90%, even more preferably about 95%, and most preferably about 97% homology, which encode an active polypeptide. For purposes of the present invention, the degree of homology between two nucleic acid sequences is determined by the Wilbur-Lipman method (Wilbur and Lipman, 1983, *Proceedings of the National Academy of Science USA* 80: 726-730) using the LASERGENE™ MEGALIGN™ software (DNASTAR, Inc., Madison, WI) with an identity table and the following multiple alignment parameters: Gap penalty of 10, and gap length penalty of 10. Pairwise alignment parameters were Ktuple=3, gap penalty=3, and windows=20.

Modification of a nucleic acid sequence encoding a cyclohexadepsipeptide synthetase of the present invention may be necessary for the synthesis of polypeptides substantially similar to the cyclohexadepsipeptide synthetase. The term "substantially similar" to the

014627588-011300
cyclohexadepsipeptide synthetase refers to non-naturally occurring forms of the enzyme. These polypeptides may differ in some engineered way from the cyclohexadepsipeptide synthetase isolated from its native source, e.g., variants of the cyclohexadepsipeptide synthetase that differ in specific activity, thermostability, pH optimum, or the like. The
5 variant sequence may be constructed on the basis of the nucleic acid sequence presented as the polypeptide encoding part of SEQ ID NO:1, e.g., a subsequence thereof, and/or by introduction of nucleotide substitutions that do not give rise to another amino acid sequence of the polypeptide encoded by the nucleic acid sequence, but which correspond to the codon
10 usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitution, see, e.g., Ford *et al.*, 1991, *Protein Expression and Purification* 2: 95-107.

It will be apparent to those skilled in the art that such substitutions can be made outside the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the cyclohexadepsipeptide synthetase encoded by the isolated nucleic acid sequence of the invention, and therefore preferably not subject to substitution, may be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (see, e.g., Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, mutations are introduced
20 at every positively charged residue in the molecule, and the resultant mutant molecules are tested for cyclohexadepsipeptide synthetase activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional structure as determined by such techniques as nuclear magnetic resonance analysis, crystallography or photoaffinity labelling (see, e.g.,
25 de Vos *et al.*, 1992, *Science* 255: 306-312; Smith *et al.*, 1992, *Journal of Molecular Biology* 224: 899-904; Wlodaver *et al.*, 1992, *FEBS Letters* 309: 59-64).

The present invention also relates to isolated nucleic acid sequences encoding a cyclohexadepsipeptide synthetase of the present invention, which hybridize under low stringency conditions, more preferably medium stringency conditions, even more preferably
30 high stringency conditions, and most preferably very high stringency conditions as defined herein with a nucleic acid probe that hybridizes under the same conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence

of (i) or (ii), or (iv) a complementary strand of (i), (ii), or (iii); or allelic variants thereof (Sambrook *et al.*, 1989, *supra*), as defined herein.

The present invention also relates to isolated nucleic acid sequences produced by (a) hybridizing a DNA under low, medium, high, or very high stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii), or (iv) a complementary strand of (i), (ii), or (iii); and (b) isolating the nucleic acid sequence. The subsequence is preferably a sequence of at least 100 nucleotides such as a sequence that encodes a polypeptide fragment, which has cyclohexadepsipeptide synthetase activity.

The present invention further relates to methods for producing a mutant nucleic acid sequence, comprising introducing at least one mutation into the mature polypeptide coding sequence of SEQ ID NO:1 or a subsequence thereof, wherein the mutant nucleic acid sequence encodes a polypeptide which consists of the mature polypeptide contained within SEQ ID NO:2 or a fragment thereof that has cyclohexadepsipeptide synthetase activity.

The introduction of a mutation into the nucleic acid sequence to exchange one nucleotide for another nucleotide may be accomplished by site-directed mutagenesis using any of the methods known in the art. Particularly useful is the procedure which utilizes a supercoiled, double stranded DNA vector with an insert of interest and two synthetic primers containing the desired mutation. The oligonucleotide primers, each complementary to opposite strands of the vector, extend during temperature cycling by means of *Pfu* DNA polymerase. On incorporation of the primers, a mutated plasmid containing staggered nicks is generated. Following temperature cycling, the product is treated with *DpnI* which is specific for methylated and hemimethylated DNA to digest the parental DNA template and to select for mutation-containing synthesized DNA. Other procedures known in the art may also be used.

The present invention also relates to nucleic acid constructs, recombinant expression vectors, and host cells containing the nucleic acid sequence of SEQ ID NO:1, subsequences or homologues thereof, for expression of the sequences. The constructs and vectors may be constructed as described herein. The host cell may be any cell suitable for the expression of the nucleic acid sequence.

The host cell may be a eukaryote, such as a mammalian cell, an insect cell, a plant cell or a fungal cell. Useful mammalian cells include Chinese hamster ovary (CHO) cells, HeLa

cells, baby hamster kidney (BHK) cells, COS cells, or any number of other immortalized cell lines available, e.g., from the American Type Culture Collection.

In a preferred embodiment, the host cell is a fungal cell. In a more preferred embodiment, the fungal host cell is a yeast cell or a filamentous fungal cell.

In an even more preferred embodiment, the yeast host cell is a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell.

In a most preferred embodiment, the yeast host cell is a *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis* or *Saccharomyces oviformis* cell. In another most preferred embodiment, the yeast host cell is a *Kluyveromyces lactis* cell.

In another most preferred embodiment, the yeast host cell is a *Yarrowia lipolytica* cell.

In another even more preferred embodiment, the filamentous fungal host cell is an *Acremonium*, *Aspergillus*, *Fusarium*, *Humicola*, *Mucor*, *Myceliophthora*, *Neurospora*, *Penicillium*, *Thielavia*, *Tolypocladium*, or *Trichoderma* cell.

In a most preferred embodiment, the filamentous fungal host cell is an *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Fusarium bacridioides*, *Fusarium crookwellense* (synonym of *Fusarium cerealis*), *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium solani*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, (e.g., *Fusarium venenatum* (Nirenberg sp. nov.)), *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophilum*, *Neurospora crassa*, *Penicillium purpurogenum*, *Thielavia terrestris*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Fusarium* host cells are described herein. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology*, Methods in Enzymology, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito et al., 1983, *Journal of Bacteriology* 153: 163; and Hinnen et al., 1978, *Proceedings of the*

The present invention also relates to methods for producing a cyclohexadepsipeptide synthetase of the present invention comprising (a) cultivating a strain, which in its wild-type form is capable of producing the cyclohexadepsipeptide synthetase, to produce a supernatant comprising the cyclohexadepsipeptide synthetase; and (b) recovering the cyclohexadepsipeptide synthetase. Preferably, the strain is of the genus *Fusarium*, and more preferably *Fusarium venenatum*.

The present invention also relates to methods for producing a cyclohexadepsipeptide synthetase of the present invention comprising (a) cultivating a host cell under conditions conducive for production of the cyclohexadepsipeptide synthetase; and (b) recovering the cyclohexadepsipeptide synthetase.

In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the cyclohexadepsipeptide synthetase using methods known in the art as described herein. The cyclohexadepsipeptide synthetase may be detected using methods known in the art specific for the enzyme (see, e.g., Visconti *et al.*, 1992, *supra*). The resulting cyclohexadepsipeptide synthetase may be recovered and purified by methods known in the art as described herein.

The present invention also relates to methods for producing cyclohexadepsipeptides and to cyclohexadepsipeptides produced by the cyclohexadepsipeptide synthetases of the present invention. The production of a cyclohexadepsipeptide may be accomplished with the isolated synthetase or by fermentation of a cell containing the gene encoding the synthetase (see, for example, Madry *et al.*, 1983, *European Journal of Applied Microbiology and Biotechnology* 17: 75-79).. The cell may be a wild-type cell or a recombinant cell. The cyclohexadepsipeptides may be isolated and purified by any of the methods known in the art. See, for example, U.S. Patent No. 5,656,464; Visconti *et al.*, 1992, *supra*.

In a preferred embodiment, the method for producing a cyclohexadepsipeptide, comprises: (a) reacting a cyclohexadepsipeptide synthetase of the present invention with D-2-hydroxyisovaleric acid, a branched chain L-amino acid, S-adenosylmethionine, and ATP; and (b) isolating the cyclohexadepsipeptide from the reaction.

In another preferred embodiment, the method for producing a cyclohexadepsipeptide, comprises: (a) cultivating a cell under conditions suitable for the production of the cyclohexadepsipeptide, wherein the cell comprises a nucleic acid sequence encoding (i) a

0948788-011300

cyclohexadepsipeptide synthetase having an amino acid sequence which has at least 65% identity with the mature polypeptide contained within SEQ ID NO:2; (ii) a cyclohexadepsipeptide synthetase which is encoded by a nucleic acid sequence which hybridizes under medium stringency conditions with the nucleic acid sequence of SEQ ID NO:1 or its complementary strand, or a subsequence of SEQ ID NO:1 of at least 100 nucleotides; (iii) an allelic variant of (a) or (b); or (iv) a fragment of (a), (b), or (c) that has cyclohexadepsipeptide synthetase activity; and (b) isolating the cyclohexadepsipeptide from the reaction.

The present invention is further described by the following examples which should not be construed as limiting the scope of the invention.

Examples

Strains

Fusarium venenatum strain ATCC 20334 was used as the source of genomic DNA for these experiments. Genomic DNA libraries were constructed using the λ ZipLox cloning system (Life Technologies, Gaithersburg, MD) with *E. coli* Y1090ZL as a host for plating and purification of recombinant bacteriophage and *E. coli* DH10Bzip for excision of recombinant pZL1-derivatives. *Fusarium torulosum* R-5690 (Fusarium Research Center, Penn State University, State College, PA) and *Aspergillus niger* Bo-1 (Novo Nordisk A/S, Bagsvaerd, Denmark) were used as sources of control DNAs for hybridization experiments. The *tri5*-deleted *Fusarium venenatum* strain LyMC1A (WO 99/60137) was used as the recipient for transformation experiments. *Escherichia coli* TOP10 (Invitrogen Corp., Carlsbad, CA) and *E. coli* DH5-alpha strains (Gibco-BRL Life Technologies, Bethesda, MD) were used for vector construction and routine plasmid propagation.

Media

RA sporulation medium was composed per liter of 50 g of succinic acid (disodium salt), 20 ml of 50X Vogels salts, 12.1 g of NaNO_3 , and 1 g of glucose.

50X Vogels Salts was composed per liter of 125 g of sodium citrate, 250 g of KH_2PO_4 , 10 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (predissolved in 20 ml water), and 5 ml of 200X Vogels trace elements. (Each ingredient was dissolved completely before addition of

the next one). Filter sterilized.

200 X Vogels Trace Elements was composed per 100 ml of 5g of citric acid·1H₂O, 5 g of ZnSO₄·7H₂O, 1 g of Fe(NH₄)₂(SO₄)₂·6H₂O, 0.25 g of CuSO₄·5H₂O, 0.05 g of MnSO₄·1H₂O, 0.05 g of H₃BO₃, and 0.05 g of Na₂MoO₄·2H₂O.

Fluoroacetamide agar (FA) was composed per liter of 12g of sodium acetate, 2 g of sodium chloride, 0.5 g of MgSO₄, 3 g of KH₂PO₄, 0.3 g of urea, 2 g of fluoroacetamide, 1 ml of Vogels salts, and 15 g of Noble agar (pH 6.1).

Cove medium was composed per liter of 342.3 g of sucrose, 20 ml of 50X Cove salt solution, 10 ml of 1 M acetamide, 10 ml of 1.5 M CsCl, and 25 g of Noble agar.

50X Cove Salts was composed per liter of 26g of KCl, 26 g of MgSO₄·7H₂O, 76g of KH₂PO₄, and 50 ml of 20X Cove trace elements.

20X Cove trace elements was composed per liter of 0.04 g of Na₂B₄O₇·10H₂O, 0.4 g of CuSO₄·5H₂O, 1.2 g of FeSO₄·7H₂O, 0.7 g of MnSO₄·H₂O, 0.8 g of Na₂MoO₄·2H₂O, and 10 g of ZnSO₄·7H₂O.

Example 1: Genomic DNA Extraction of *Fusarium venenatum*, *Fusarium torulosum*, and *Aspergillus niger*

Fusarium venenatum, *Fusarium torulosum*, and *Aspergillus niger* were each grown for 24-36 hours at 28°C and 150 rpm in 25 ml of YEG medium composed per liter of 5 g of yeast extract and 20 g of glucose. Mycelia were then collected by filtration through Miracloth (Calbiochem, La Jolla, CA) and washed once with 25 ml of 10 mM Tris-1 mM EDTA (TE) buffer. Excess buffer was drained from the mycelia which were subsequently frozen in liquid nitrogen. The frozen mycelia were ground to a fine powder in an electric coffee grinder, and each powder was added to 20 ml of TE buffer and 5 ml of 20% w/v sodium dodecylsulfate (SDS) in a disposable plastic centrifuge tube. The mixtures were gently inverted several times to ensure mixing, and extracted twice with an equal volume of phenol:chloroform:isoamyl alcohol (25:24:1 v/v/v). Sodium acetate (3 M solution) was added to give a final concentration of 0.3 M and the nucleic acids were precipitated with 2.5 volumes of ice cold ethanol. The tubes were centrifuged at 15,000 x g for 30 minutes and the pellets were allowed to air dry for 30 minutes before resuspension in 0.5 ml of TE buffer. DNase-free ribonuclease A was added to a concentration of 100 µg/ml and the mixtures were incubated at 37°C for 30 minutes. Proteinase K (200 µg/ml) was then added and the mixtures

5 were incubated an additional hour at 37°C. Finally, the mixtures were extracted twice with phenol:chloroform:isoamyl alcohol (25:24:1 v/v/v) before precipitating the DNA with sodium acetate and ethanol according to standard procedures. The DNA pellets were dried under vacuum, resuspended in TE buffer, and stored at 4°C.

Example 2: Hybridization Experiments

10 The genomic DNA preparations described in Example 1 were tested for the presence of cyclohexadepsipeptide synthetase gene sequences using Southern hybridization. Aliquots of the DNA were digested with *Bam*HI or *Bam*HI plus *Xba*I and fractionated by agarose gel electrophoresis. The DNA in the gel was blotted to a Hybond N+™ membrane filter (Amersham Corporation, Arlington Heights, IL) according to the method of Davis *et al.* (1980, *Advanced Bacterial Genetics, A Manual for Genetic Engineering*, Cold Spring Harbor Press, Cold Spring Harbor, NY), and probed with a radiolabeled fragment encoding the 5' portion of the *Fusarium torulosum esyn1* gene (obtained from Dr. Thomas Hohn, USDA, Peoria, IL) under low, medium, and high stringency hybridization conditions at 45°C as described herein. The cyclohexadepsipeptide synthetase-specific probe fragment from *Fusarium torulosum* was radiolabeled by nick translation (Sambrook *et al.*, 1989, *Molecular Cloning. A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, NY) with α [³²P]dCTP (Amersham, Arlington Heights, IL), denatured by adding NaOH to a final concentration of 0.1 M, and added to the hybridization buffer at an activity of approximately 1 x 10⁶ cpm per ml. Following the hybridization, the filters were washed once in 0.2X SSPE with 0.1% SDS at 45°C followed by two washes in 0.2X SSPE (no SDS) at the same temperature. The filters were allowed to dry on paper towels for 15 minutes, then wrapped in Saran-wrap™ and exposed to X-ray film overnight at -70°C with intensifying screens 25 (Kodak, Rochester, NY).

30 Southern hybridization analysis showed that cyclohexadepsipeptide synthetase-specific DNA sequences could be detected in the genome of *Fusarium venenatum* with the *Fusarium torulosum esyn1* probe only under conditions of low and medium stringency. The positive control DNA from *Fusarium torulosum* gave strong hybridization signals under all conditions, and negative control DNA from *Aspergillus niger* failed to hybridize under all conditions tested. These results suggested that *Fusarium venenatum* contained genomic DNA sequences homologous to the *Fusarium torulosum* enniatin synthetase gene.

Example 3: Genomic DNA Library Construction and Screening

Genomic libraries of *Fusarium venenatum* were constructed in λ ZipLox according to the manufacturer's instructions (Life Technologies, Gaithersburg, MD). *Fusarium venenatum* genomic DNA was partially digested with *Tsp*509I and size-fractionated on 1% agarose gels. DNA fragments migrating in the size range 3-7 kb were excised and eluted from the agarose gel slices using Prep-a-Gene reagents (BioRad, Hercules, CA). The eluted DNA fragments were ligated with *Eco*RI-cleaved and dephosphorylated λ ZipLox vector arms (Life Technologies, Gaithersburg, MD), and the ligation mixtures were packaged using commercial packaging extracts (Stratagene, La Jolla, CA). The packaged DNA libraries were plated and amplified in *E. coli* Y1090ZL cells.

Approximately 50,000 plaques from the library were screened by plaque-hybridization (Davis *et al.*, 1980, *supra*) with the radiolabeled probe fragment of the *Fusarium torulosum esyn1* gene using the low stringency conditions described in Example 2. Plaques, which gave hybridization signals, were purified twice in *E. coli* Y1090ZL cells, and the individual clones were subsequently excised from the λ ZipLox vector as pZL1-derivatives (D'Alessio *et al.*, 1992, *Focus*® 14: 7). Chromosome "walking" to obtain adjacent DNA sequences was done using homologous *Fusarium venenatum* probes at high stringency.

Four plaques were identified that hybridized strongly to the *Fusarium torulosum esyn1* gene probe, and each of the potential clones was subsequently excised from the λ ZipLox vector as a pZL1-derivative (D'Alessio *et al.*, 1992, *supra*). Plasmid DNA was isolated from the clones by passage through *E. coli* DH10B cells using standard methods. The sizes of the cloned inserts were determined by agarose gel electrophoresis. The largest insert comprised a DNA segment of approximately 3 kb. The clone was designated *E. coli* DH10B pZL-ESA.

Example 4: Cloning and Analysis of a *Fusarium venenatum* Cyclohexadepsipeptide Synthetase Gene

DNA sequencing of the DNA segment of approximately 3 kb was performed with an Applied Biosystems Model 377 XL Automated DNA Sequencer using dye-terminator chemistry. Contiguous sequences were generated using a transposon insertion strategy

(Primer Island Transposition Kit, Perkin-Elmer/Applied Biosystems, Inc., Foster City, CA). The entire cloned region was sequenced to an average redundancy of 6.9.

Nucleotide sequencing revealed that the 3 kb segment contained an open reading frame encoding at least 900 amino acids. However, this fragment (designated Fragment A, pZL-ESA) did not encode the entire gene product. Consequently, the library was re-screened using a probe comprising the 3'-portion of Fragment A (ca. 1 kb *Hind*III fragment). Several clones were subsequently identified and analyzed by restriction mapping. The largest of these secondary clones contained a genomic DNA insert of about 4.6 kb (designated Fragment B, pZL-ESB). The clone was designated *E. coli* DH10B pZL-ESB.

Nucleotide sequence examination of Fragment B extended the open reading frame of Fragment A by amino acids 777 through 2311. However, this sequence did not reach the stop codon of the open reading frame, thereby necessitating isolation of a third genomic segment. The third genomic clone was isolated by re-screening the genomic library with a PCR-amplified probe derived from Fragment B. Two PCR primers shown below were used to amplify a 586 bp probe segment used for screening the library.

5'-daATTGATTTCGCTTGAAAGTCGAT-3' (SEQ ID NO:3)

5'-dCTTGAGAGTTACGTTGGTCTTGAAC-3' (SEQ ID NO:4)

The amplification reaction (100 μ l) contained the following components: 0.2 μ g of pZL-ESB DNA, 48.4 pmol of the forward primer, 48.4 pmol of the reverse primer, 1 mM each of dATP, dCTP, dGTP, and dTTP, 1 x *Taq* polymerase buffer, and 2.5 U of *Taq* polymerase (Perkin-Elmer Corp., Branchburg, NJ). The reaction was incubated in an Ericomp Twin Block System Easy Cycler programmed for 1 cycle at 95°C for 5 minutes followed by 30 cycles each at 95°C for 1 minute, 55°C for 1 minute and 72°C for 2 minutes.

The reaction was electrophoresed on an agarose gel, and the expected product of 586 bp was obtained. The reaction was run on a preparative gel, a gel slice containing the desired product was excised, and DNA was isolated from the gel using a Qiaquick Gel Extraction Kit (Qiagen, Chatsworth, CA).

From seven clones that were identified with this probe, the largest (Fragment C, pZL-ESC) contained a 5.5 kb insert. Subsequent DNA sequence analysis revealed that Fragment C encoded amino acids 1617 through 3129, a potential stop codon, and 1553 bp of 3'-flanking DNA. The clone was designated *E. coli* DH10B pZL-ESC. The entire DNA sequence of the cyclohexadepsipeptide synthetase gene was assembled from the three

overlapping clones (Fragments A, B, and C). A transposon insertion strategy allowed for rapid sequencing to high redundancy.

The complete DNA sequence and deduced amino acid sequence are shown in Figure 1. The DNA sequence of the cyclohexadepsipeptide synthetase gene (SEQ ID NO:1) was determined to an average redundancy of 6.9. The cyclohexadepsipeptide synthetase gene contained a lengthy open reading frame of 9387 bp with no introns, encoding a polypeptide of 3129 amino acids (MW = 346,852).

The deduced amino acid sequence (SEQ ID NO:2) of the cyclohexadepsipeptide synthetase gene product shared approximately 59% identity to the enniatin synthetase of *Fusarium scirpi* (Haese *et al.*, 1993, *Mol. Microbiol.* 7: 905-914; DNA sequence listed in EMBL database under accession number Z18755). Percent identity was determined by the Clustal method (Higgins, 1989, *CABIOS* 5: 151-153) using the LASERGENE™ MEGALIGN™ software (DNASTAR, Inc., Madison, WI) with an identity table and the following multiple alignment parameters: Gap penalty of 10 and gap length penalty of 10. Pairwise alignment parameters were Ktuple=1, gap penalty=3, windows=5, and diagonals=5.

Example 5: pΔES-*amdS* Construction

The construction of the *dps1* deletion vectors pΔES-*amdS*1 and pΔES-*amdS*2 is shown in Figure 2. Briefly, a 0.2 kb DNA segment comprising a portion of the *dps1* coding region was removed from plasmid pZL-ESA (designated as fragment A) by digestion with *Stu*I and *Nru*I restriction endonucleases. Both of these enzymes generate blunt-ended DNA fragments. The digested pESA vector was treated with calf intestine alkaline phosphatase (Boehringer-Mannheim Biochemicals, Indianapolis, IN) to prevent self-ligation. Lastly, a 3.2 kb fragment encoding the *Aspergillus nidulans amdS* gene (with flanking repeat sequences derived from the *Aspergillus oryzae pyrG* gene) was obtained by digestion of pJRoy47 (WO 99/60137) with *Sma*I and *Pme*I. This *amdS* fragment was subsequently ligated with the pZL-ESA vector fragment described above to generate the deletion plasmids pΔES-*amdS*1 and pΔES-*amdS*2 (which differ only in the orientation of the *amdS* gene segment).

Example 6: Transformation of *Fusarium venenatum* LyMC1A and Preliminary Screening for *dps1* Gene Deletions

Plasmid pΔES-*amdS*1 was digested with *Spe*I, and the 5.7 kb deletion fragment

(comprising portions of the *Fusarium venenatum dps1* gene with the *Aspergillus nidulans amdS* gene and repeats replacing 0.2 kb of the *dps1* coding region) was subsequently excised and purified for use in transformation experiments. The preparation and transformation of *Fusarium venenatum* LYMC1A protoplasts was performed according to the method of Royer, 1995, *Bio/Technology* 13: 1479-1483.

Fusarium venenatum LYMC1A protoplasts were transformed with the 5.7 kb *SpeI* Δ ES-*amdS* fragment with selection on COVE plates. Fifteen transformants were obtained and single spore purified. DNA was extracted from the single spore purified transformants, generated with the *SpeI* Δ ES-*amdS* fragment, as well as from *Fusarium venenatum* LYMC1A, using the Qiagen DNeasy Plant mini kit (Qiagen, Chatsworth, CA) (with a 2 hour lytic incubation in place of 10 minutes recommended in the manufacturer's protocol). One to two micrograms of each DNA were digested for seven hours with *XhoI* or *SpeI* (10 U/ μ g DNA in 30 μ l reactions). The digests were electrophoresed on 1% agarose gels in TAE buffer, and the DNAs were transferred to Hybond N⁺ in 0.4 N NaOH. The blots were UV crosslinked and probed as described below.

Probes were prepared using the Prime-It Labeling Kit (Stratagene, La Jolla, CA) and α [³²P]-dCTP. Following labeling the probes were separated from unincorporated label using a G 50 TE Midi column (5' to 3', Boulder, CO).

Blots were prehybridized at 65°C in Rapid Hyb Buffer (Amersham, Arlington Heights, IL) for 45 minutes. Denatured probes were added to the Rapid Hyb solution and hybridizations were done overnight at 65°C. Following hybridization the blots were washed once at room temperature in 2X SSC for 5 minutes and in 0.2X SSC, 0.1% SDS at 65°C for 5 minutes twice. The washed blots were washed in 2X SSC at room temperature for 5 minutes.

Southern blots of *XhoI* and *SpeI*-digested genomic DNA were probed twice. First, they were probed with an 800 bp *NsiI/SpeI* fragment of p Δ ES-*amdS1*. Four of the fifteen transformants had the 5.2 kb band (*XhoI* digested DNA) and the 5.7 kb band (*SpeI* digested DNA) expected for a gene replacement when probed with the 800 bp *NsiI/SpeI* fragment. Most of the other transformants had the 2.2 kb band (*XhoI* digested DNA) or the 2.7 kb (*SpeI* digested DNA) wild-type bands, and additional bands, most likely corresponding to ectopic integration of the transforming DNA.

Secondly, the same Southern blots were probed with *HindIII*-linearized pDSY176, a plasmid containing the 0.2 kb *StuI/NruI* portion of the *dps1* coding region. pDSY176 was

constructed as follows: pZL-ESA was digested with *StuI*/*NruI*, and the 0.2 kb fragment was isolated by preparative electrophoresis. The isolated fragment was cloned into pZERO-Blunt (Invitrogen Corp., Carlsbad, CA) according to the manufacturer's instructions to produce pDSY176. Hybridization analysis using this second probe (pDSY176) confirmed that none of the four putative deleted strains (*Fusarium venenatum* Δ ES 4, 6, 8, and 10) contained the 0.2 kb region of the *dpsI* gene which had been deleted.

Example 7: Removal of the *amdS* Gene

Two of the transformants confirmed as being deleted, Δ ES4A and Δ ES8A, were sporulated in 500 ml of RA medium, in 2 liter Fernbach flasks. These were inoculated with 12 mycelial plugs cut from Cove plates and incubated at 24°C, 150 rpm for 53 hours. After this time spores were harvested through sterile Miracloth (Calbiochem, San Diego, CA) centrifuged for 30 minutes at 7,000 rpm in a Sorval GS3 rotor and washed three times with sterile water. Freshly harvested spores were plated at 10^4 , 10^5 and 10^6 per plate (five plates at each concentration) of FA. Colonies which grew on these plates were picked to FA and Cove plates.

Numerous colonies were obtained on FA plates (primarily from plates seeded with 10^5 and 10^6 spores/plate) which, on subculturing, grew well on FA but only sparsely on Cove. Thirty two Δ ES4A-derived colonies (designated as *Fusarium venenatum* WTY700-3-4) and 128 Δ ES8A-derived colonies (designated as *Fusarium venenatum* WTY700-3-8) all had an *amdS*-minus phenotype.

DNA was extracted from five *Fusarium venenatum* WTY700-3-4 isolates (WTY700-3-4a through 4e) and ten *Fusarium venenatum* WTY700-3-8 strains (WTY700-3-8a through 8j) as well as LyMC1A, Δ ES4A and Δ ES8A using the Qiagen DNeasy Plant Mini Kit (with a 2 hour lytic incubation in place of 10 min recommended in the manufacturer's protocol). One microgram of each DNA was digested overnight with *XhoI* and *SpeI* (20 U/ μ g DNA in 50 μ l reactions). The digests were concentrated to 10 μ l and run on 1% agarose gels in TBE. DNAs were transferred to Hybond N+ membranes using the manufacturer's protocol, and were then probed with the probes described in Example 6.

Hybridization analysis of genomic DNAs extracted from these strains (using the 800 bp *NsiI*-*SpeI* fragment of Δ ES-*amdS1* and *HindIII*-digested pDSY176 (which contains the 200 bp *StuI*-*NruI* portion of the *dpsI* open reading frame) revealed that none of the WTY700-

3-4 isolates had lost the *amdS* gene, but 50% (5/10) of the *Fusarium venenatum* WTY700-3-8 isolates showed a banding pattern consistent with removal of the *amdS* gene. Hybridization analysis, thus, confirmed that *Fusarium venenatum* WTY700-3-8a, b, c, d, and e were deleted for the 200 bp *StuI/NruI* portion of the *dpsI* open reading frame. As expected for a deletion, a hybridization signal was observed using the pDSY176 probe described above for the parent strain *Fusarium venenatum* LyMC1A but not for any of the five *Fusarium venenatum* WTY700-3-8a-e strains.

Deposit of Biological Materials

The following biological materials have been deposited under the terms of the Budapest Treaty with the Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL), Peoria, Illinois, and given the following accession numbers:

Deposit	Accession Number	Date of Deposit
<i>E. coli</i> DH10B (pZL-ESA)	NRRL B-30068	October 27, 1998
<i>E. coli</i> DH10B (pZL-ESB)	NRRL B-30069	October 27, 1998
<i>E. coli</i> DH10B (pZL-ESC)	NRRL B-30070	October 27, 1998

The strains have been deposited under conditions that assure that access to the cultures will be available during the pendency of this patent application to one determined by the Commissioner of Patents and Trademarks to be entitled thereto under 37 C.F.R. §1.14 and 35 U.S.C. §122. The deposits represent substantially pure cultures of the deposited strains. The deposits are available as required by foreign patent laws in countries wherein counterparts of the subject application, or its progeny are filed. However, it should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by governmental action.

The invention described and claimed herein is not to be limited in scope by the specific embodiments herein disclosed, since these embodiments are intended as illustrations of several aspects of the invention. Any equivalent embodiments are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the

foregoing description. Such modifications are also intended to fall within the scope of the appended claims. In the case of conflict, the present disclosure including definitions will control.

Various references are cited herein, the disclosures of which are incorporated by
5 reference in their entireties.

Claims

What is claimed is:

1. A method for producing a heterologous polypeptide, comprising:

(a) cultivating a mutant of a parent filamentous fungal cell under conditions conducive for the production of the heterologous polypeptide, wherein (i) the mutant comprises a first nucleic acid sequence encoding the heterologous polypeptide, and (ii) the mutant produces less of a cyclohexadepsipeptide than the parent filamentous fungal cell when cultured under the same conditions; and

(b) isolating the heterologous polypeptide from the cultivation medium.

2. The method of claim 1, wherein the filamentous fungal cell is an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Gibberella*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Myrothecium*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Piromyces*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, or *Trichoderma* cell.

3. The method of claim 1, wherein the filamentous fungal cell is a *Fusarium* cell.

4. The method of claim 3, wherein the *Fusarium* cell is a *Fusarium venenatum* cell.

5. The method of claim 4, wherein the *Fusarium venenatum* cell is *Fusarium venenatum* ATCC 20334.

6. The method of claim 4, wherein the *Fusarium venenatum* cell is a morphological mutant.

7. The method of claim 6, wherein the *Fusarium venenatum* cell is a morphological mutant of *Fusarium venenatum* ATCC 20334.

8. The method of any of claims 1-7, wherein the mutant cell comprises a second nucleic acid sequence which comprises a modification of at least one of the genes involved in the production of the cyclohexadepsipeptide.

9. The method of claim 8, wherein the genes are selected from the group consisting of a cyclohexadepsipeptide synthetase gene, enniatin synthetase gene, and D-hydroxyisovalerate dehydrogenase gene.

10. The method of claim 8, wherein one of the genes is a cyclohexadepsipeptide synthetase gene.

11. The method of claim 8, wherein one of the genes is an enniatin synthetase gene.

12. The method of claim 8, wherein one of the genes is a D-hydroxyisovalerate dehydrogenase gene.

13. The method of any of claims 1-12, wherein the mutant cell produces at least about 25% less of the cyclohexadepsipeptide than the parent filamentous fungal cell when cultured under identical conditions.

14. The method of any of claims 1-13, wherein the mutant cell produces no cyclohexadepsipeptide.

15. The method of any of claims 1-14, wherein the filamentous fungal cell comprises at least two copies of the first nucleic acid sequence.

16. The method of any of claims 1-15, wherein the heterologous polypeptide is a hormone, hormone variant, enzyme, receptor or portion thereof, antibody or portion thereof, or reporter.

17. The method of claim 16, wherein the enzyme is an oxidoreductase, transferase,

hydrolase, lyase, isomerase, or ligase.

18. The method of claim 17, wherein the enzyme is an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, invertase, laccase, lipase, mannosidase, mutanase, oxidase, a pectinolytic enzyme, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, or xylanase.

19. The method of any of claims 1-18, wherein the mutant cell further comprises one or more modifications of one or more third nucleic acid sequences, wherein the modification reduces or eliminates expression of the one or more third nucleic acid sequences.

20. The method of claim 19, wherein the third nucleic acid sequence encodes an enzyme selected from the group consisting of an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, invertase, laccase, lipase, mannosidase, mutanase, oxidase, pectinolytic enzyme, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, and xylanase.

21. The method of any of claims 19, wherein the third nucleic acid sequence encodes a protease.

22. A cyclohexadepsipeptide-deficient mutant of a filamentous fungal cell, comprising a first nucleic acid sequence encoding a heterologous polypeptide, wherein the mutant produces less of a cyclohexadepsipeptide than the parent filamentous fungal cell of the mutant cell when cultured under the same conditions.

23. The mutant cell of claim 22, wherein the mutant cell comprises a second nucleic acid sequence which comprises a modification of at least one of the genes involved in the

production of the cyclohexadepsipeptide.

24. The mutant cell of claim 23, wherein the genes are selected from the group consisting of a cyclohexadepsipeptide synthetase gene, enniatin synthetase gene, and D-hydroxyisovalerate dehydrogenase gene.

25. The mutant cell of claim 23, wherein one of the genes is a cyclohexadepsipeptide synthetase gene.

26. The mutant cell of claim 23, wherein one of the genes is an enniatin synthetase gene.

27. The mutant cell of claim 23, wherein one of the genes is a D-hydroxyisovalerate dehydrogenase gene.

28. The mutant cell of any of claims 22-27, wherein the cell comprises at least two copies of the first nucleic acid sequence.

29. A method for obtaining the mutant cell of any of claims 22-28, comprising:

(a) introducing into a parent filamentous fungal cell a first nucleic acid sequence encoding a heterologous polypeptide and a second nucleic acid sequence comprising a modification of at least one of the genes responsible for the production of a cyclohexadepsipeptide; and

(b) identifying the mutant from step (a), wherein the mutant produces less of the cyclohexadepsipeptide than the parent filamentous fungal cell of the mutant cell when cultured under the same conditions.

30. An isolated cyclohexadepsipeptide synthetase, selected from the group consisting of:

(a) a cyclohexadepsipeptide synthetase having an amino acid sequence which has at least 65% identity with the mature polypeptide contained within SEQ ID NO:2;

(b) a cyclohexadepsipeptide synthetase which is encoded by a nucleic acid sequence which hybridizes under medium stringency conditions with (i) the nucleic acid

sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii) of at least 100 nucleotides, or (iv) a complementary strand of (i), (ii), or (iii);

(c) an allelic variant of (a) or (b); and

(d) a fragment of (a), (b), or (c) that has cyclohexadepsipeptide synthetase activity.

31. The cyclohexadepsipeptide synthetase of claim 30, having an amino acid sequence which has at least 65% identity with the mature polypeptide contained within SEQ ID NO:2.

32. The cyclohexadepsipeptide synthetase of claim 31, having an amino acid sequence which has at least 70% identity with the mature polypeptide contained within SEQ ID NO:2.

33. The cyclohexadepsipeptide synthetase of claim 32, having an amino acid sequence which has at least 80% identity with the mature polypeptide contained within SEQ ID NO:2.

34. The cyclohexadepsipeptide synthetase of claim 33, having an amino acid sequence which has at least 90% identity with the mature polypeptide contained within SEQ ID NO:2.

35. The cyclohexadepsipeptide synthetase of claim 34, having an amino acid sequence which has at least 95% identity with the mature polypeptide contained within SEQ ID NO:2.

36. The cyclohexadepsipeptide synthetase of claim 30, comprising the amino acid sequence of SEQ ID NO:2.

37. The cyclohexadepsipeptide synthetase of claim 30, consisting of the amino acid sequence of SEQ ID NO:2 or a fragment thereof.

38. The cyclohexadepsipeptide synthetase of claim 37, consisting of the amino acid sequence of SEQ ID NO:2.

39. The cyclohexadepsipeptide synthetase of claim 38, which consists of the mature

polypeptide contained within SEQ ID NO:2.

40. The cyclohexadepsipeptide synthetase of any of claims 30-39, which is obtained from a *Fusarium* strain.

41. The cyclohexadepsipeptide synthetase of claim 40, which is obtained from *Fusarium venenatum* strain.

42. The cyclohexadepsipeptide synthetase of claim 30, which is encoded by a nucleic acid sequence which hybridizes under medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii) of at least 100 nucleotides, or (iv) a complementary strand of (i), (ii), or (iii);.

43. The cyclohexadepsipeptide synthetase of claim 42, which is encoded by a nucleic acid sequence which hybridizes under medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, or (iii) a complementary strand of (i) or (ii).

44. The cyclohexadepsipeptide synthetase of claim 42 or 43, which is obtained from a *Fusarium* strain.

45. The cyclohexadepsipeptide synthetase of claim 44, which is obtained from *Fusarium venenatum* strain.

46. The cyclohexadepsipeptide synthetase of claim 30, which is encoded by a nucleic acid sequence which hybridizes under high stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii) of at least 100 nucleotides, or (iv) a complementary strand of (i), (ii), or (iii).

47. The cyclohexadepsipeptide synthetase of claim 46, which is encoded by a nucleic acid sequence which hybridizes under high stringency conditions with (i) the nucleic acid

sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, or (iii) a complementary strand of (i), (ii), or (iii);

48. The cyclohexadepsipeptide synthetase of claim 46 or 47, which is obtained from a *Fusarium* strain.

49. The cyclohexadepsipeptide synthetase of claim 48, which is obtained from *Fusarium venenatum* strain.

50. The cyclohexadepsipeptide synthetase of claim 30, which is encoded by the nucleic acid sequences contained in plasmid pZL-ESA, which is contained in *Escherichia coli* NRRL B-30068, plasmid pZL-ESB, which is contained in *Escherichia coli* NRRL B-30069, and plasmid pZL-ESC, which is contained in *Escherichia coli* NRRL B-30070.

51. The cyclohexadepsipeptide synthetase of any of claims 30-50 which has at least 20% of the enzyme activity of SEQ ID NO:2.

52. A cyclohexadepsipeptide synthetase having the same enzyme activity as the cyclohexadepsipeptide synthetase of any of claims 30-51.

53. An isolated nucleic acid sequence comprising a nucleic acid sequence which encodes the cyclohexadepsipeptide synthetase of any of claims 30-52.

54. An isolated nucleic acid sequence comprising a nucleic acid sequence having at least one mutation in the mature cyclohexadepsipeptide synthetase coding sequence of SEQ ID NO:1, in which the mutant nucleic acid sequence encodes a cyclohexadepsipeptide synthetase consisting of the mature polypeptide contained within SEQ ID NO:2.

55. An isolated nucleic acid sequence produced by (a) hybridizing a DNA under medium stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii) of at least 100 nucleotides, or (iv)

a complementary strand of (i), (ii), or (iii); and (b) isolating the nucleic acid sequence.

56. The isolated nucleic acid sequence of claim 55 produced by (a) hybridizing a DNA under high stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1, (ii) the cDNA sequence of SEQ ID NO:1, (iii) a subsequence of (i) or (ii) of at least 100 nucleotides, or (iv) a complementary strand of (i), (ii), or (iii); and (b) isolating the nucleic acid sequence.

57. A nucleic acid construct comprising the nucleic acid sequence of claim 53 operably linked to one or more control sequences that direct the production of the cyclohexadepsipeptide synthetase in a suitable expression host.

58. A recombinant expression vector comprising the nucleic acid construct of claim 57.

59. A recombinant host cell comprising the nucleic acid construct of claim 57.

60. A method for producing a mutant nucleic acid sequence, comprising (a) introducing at least one mutation into the mature cyclohexadepsipeptide synthetase coding sequence of SEQ ID NO:1, wherein the mutant nucleic acid sequence encodes a cyclohexadepsipeptide synthetase consisting of the mature polypeptide contained within SEQ ID NO:2; and (b) recovering the mutant nucleic acid sequence.

61. A mutant nucleic acid sequence produced by the method of claim 60.

62. A method for producing a cyclohexadepsipeptide synthetase, comprising (a) cultivating a strain comprising the mutant nucleic acid sequence of claim 61 encoding the cyclohexadepsipeptide synthetase to produce a supernatant comprising the cyclohexadepsipeptide synthetase; and (b) recovering the cyclohexadepsipeptide synthetase.

63. A method for producing the cyclohexadepsipeptide synthetase of any of claims 30-52 comprising (a) cultivating a strain to produce a supernatant comprising the cyclohexadepsipeptide synthetase; and (b) recovering the cyclohexadepsipeptide synthetase.

64. A method for producing the cyclohexadepsipeptide synthetase of any of claims 30-52 comprising (a) cultivating a host cell comprising a nucleic acid construct comprising a nucleic acid sequence encoding the cyclohexadepsipeptide synthetase under conditions
5 suitable for production of the cyclohexadepsipeptide synthetase; and (b) recovering the cyclohexadepsipeptide synthetase.

65. A method for producing a cyclohexadepsipeptide synthetase comprising (a) cultivating a host cell under conditions conducive for production of the
10 cyclohexadepsipeptide synthetase, wherein the host cell comprises a mutant nucleic acid sequence having at least one mutation in the mature cyclohexadepsipeptide synthetase coding sequence of SEQ ID NO:1, wherein the mutant nucleic acid sequence encodes a cyclohexadepsipeptide synthetase consisting of the mature polypeptide contained within SEQ ID NO:2, and (b) recovering the cyclohexadepsipeptide synthetase.

66. A method for producing a cyclohexadepsipeptide, comprising:

(a) reacting the cyclohexadepsipeptide synthetase of any of claims 30-52 with D-2-hydroxyisovaleric acid, a branched chain L-amino acid, S-adenosylmethionine, and ATP;
and

(b) isolating the cyclohexadepsipeptide from the reaction.

67. A cyclohexadepsipeptide produced by the method of claim 66.

68. A method for producing a cyclohexadepsipeptide, comprising:

(a) cultivating a cell under conditions suitable for the production of the
25 cyclohexadepsipeptide, wherein the cell comprises a nucleic acid sequence encoding (i) a cyclohexadepsipeptide synthetase having an amino acid sequence which has at least 65% identity with the mature polypeptide contained within SEQ ID NO:2; (ii) a cyclohexadepsipeptide synthetase which is encoded by a nucleic acid sequence which
30 hybridizes under medium stringency conditions with the nucleic acid sequence of SEQ ID NO:1 or its complementary strand, or a subsequence of SEQ ID NO:1 of at least 100

Abstract

5 The present invention relates to methods for producing a heterologous polypeptide, comprising: (a) cultivating a mutant of a parent filamentous fungal cell under conditions conducive for the production of the heterologous polypeptide, wherein (i) the mutant cell comprises a nucleic acid sequence encoding the heterologous polypeptide and (ii) the mutant produces less of the cyclohexadepsipeptide than the parent filamentous fungal cell when cultured under the same conditions; and (b) isolating the heterologous polypeptide from the cultivation medium. The present invention also relates to mutants of filamentous fungal cells and methods for obtaining the mutant cells. The present invention also relates to isolated cyclohexadepsipeptide synthetases and isolated nucleic acid sequences encoding the cyclohexadepsipeptide synthetases. The present invention also relates to nucleic acid constructs, vectors, and host cells comprising the nucleic acid sequences as well as methods for producing the cyclohexadepsipeptide synthetases. The present invention further relates to cyclohexadepsipeptides produced by the cyclohexadepsipeptide synthetases.

10

AATTAGATTCCACTAGTAGCCATTGTAGATCAAGGCCAAGATATGACAAACCCATAAGTAACGGCGATCCTGTCTCAT 80
 GTATCCAAAATAAGAGACACGGGATATTCTACTTTTCAGATCTTTCTTCAAATCTCTCCCTCGAGAAGCTACTTGGGA 160
 TGAATTGAGTCTTCTGGCTCAGATAGATATCTACTGTATCTCCGATAGACTTTTGCCTGGTAGCATTAACGTTCTCTTA 240
 TATTCTAATTGTAATTCATTACATATGGAATATTCTACTGCTCGATGGTAGGCAAGACCTTGCCACCTACACCA 320
 MEYLTATADV DGRQLPLPTP
 GCTTCGTTTTGTAGTCATGGAGATAGTCCCTCCATATAGCTTTACGAGCAACTCTTCCATCTCTATGTCTGATTCGAG 400
 ASFC SHGDSPLN SSVYEQ L F H L Y G L D S S
 TCGCATCGAAGCTATCAAAACCATGCACACCTTTCCAGCTTGACATGATGCACTGCAATGCTTTGGATAAGCAAGTCTGCTA 480
 RIEAIK PCTPQLDMIDCNALDKQSA
 TCGGCCATGCGGTGATGATGTCCAAACCGACATTCACATCTCTCGTTTCGCGCTTGCCTGGAGGAGAGATCGTCAACCAA 560
 IGHAVYDVP TDI D I S R F A L A W K E I V N Q
 ACCCCAGCCTTGCGAGCGCTTTGCTTTCACCTCGGACTCTGGAAGACTTCTCAAGTCATCCTAAAGATAGCTTTGTCTT 640
 TPALRAFAFAFTSDSGKTSQVILKDSFVF
 CTCATGGATGTCTGCTTCTTCGAGCTCCCCAGATGAAGTGTTCGGGATGAAGCTGCGCTGCTGCATCGGCGCAC 720
 S W M C W S S S P D E V V R D E A A A A S G P
 GCTGCAACCGCTTCGTTCTACTTTGAAGACATCGACACGAGAATGTACGTGGTTTGGACCTTCAGTCATGCAATTGGTA 800
 RCNRFVLLLEDMDMTKKQLVWTFSLALV
 GAGTCACTTTCACACACGCGCTCTGAGCGGTGTTTCGCGGTACAAAGCATGAGAAGACACACATCGGCTGAGAC 880
 D V T F Q Q R V L S R V F A A Y K H E K D T H R P E T
 ACCGAGTCATGTATGCCACTGACATGACTCTCAGTCAGTCTCCGTTGTTCCATGAGCTGCGAGGACAATGCCGTAT 960
 P E S S D A T D T D S Q S V S V S M S C E D N A V
 CGGCGACTCATTTCTGGCAAACTCACCTTAAGATCTCAATGCTCGCTCTCCCTCACCTGTCTGACCACCTGATGGTG 1040
 S A T H F W Q T H L N D L N A S V F P H L S D H L M V
 CCGAACCCACTACACACGACATCGTATACATTCCTCTTTCACAGAAGCACTACCAATTCGCGCATCTCGCC 1120
 P N P T T A E H R I T F P L S Q K A L S N S A I C R
 TACTGCACTCAATACTCTCTCGCGCTACACTCATCTGACGAGGCTGTGTTGGTGGCTGAAGTCACTGACGCAATCTAC 1200
 T A L S I L S R Y T H S D E A L F G A V T E O S L
 CATTTGACAAACACTATCTTGAGATGGTAGTACCAACAGTTGACCCCTCTCGTGTACACTGCCCAATCAATCTTCGT 1280
 P F D K H Y L A D G T Y Q T V A P P L R V H C Q S N L R

Fig. 1A

GCATCAGATGTCATGGATGCAATCTCTTCTTACGATGATCGCTTGGTCTATCTCGCCCCATTTGGCTTCGCGACATCG 1360
 A A S D V M D A I S S Y D D R L G H L A P F G L R D I R
 CAACCTGGTGAATAAGGCTTCGCCGCTCGATTTCCTCCAAATCTGTTTACTCTACCGAGTGGACGCCACGTAACAATG 1440
 N T G D N G S A C G D F Q T V L L V T D G S H V N N
 GTATCAACGGTTTCTCCAAACAGATAACAGAGTCAAGCCATTTCATGCTTGCAACAACCGTGCCCTCTCTTGCACGTG 1520
 G I N F L Q Q I T E S H F M P C N R A L L L H C
 CAGATGGAAGATGCGGAGCTCTGCTGTGCTACTATGACCAAAATTTATCGATTTCGATTTCAGACAACGCGTCTCTG 1600
 Q M E S S G A L L V A Y Y D H N V I D S L Q T T R L L
 ACAGCAGTTTGGTCACTGATCAAGTGTTCGAAAGTCCTTAGACCTGAGCTCTATGGCTGAGGTCAACTTGTGACTG 1680
 Q Q F G H L I K C L Q S P L D L S M A E V N L M T
 AGTATGACAGACAGAGATTGAGATTGGAATCGAACCGTTAGAGTACAGGATACCCGTGATCCACCATGATGTTG 1760
 E Y D R A E I E S W N S Q P L E V Q D T L I H E M L
 AAAGCTGTTTCTCAITTCGCCCAACCAACGGCCATCCAAAGGTGGATGGAGACTGGACCTATTTCGGAGCTCGACAATGT 1840
 K A V S H S P T K T A I Q A W D G D W T Y S E L D N V
 TTCGTCAAGACTCGCTGCCATATCAAGTCACTTGGCTTAGAGCTCAGCAAGCCATATTCCAGTCTACTTTGAGAAGT 1920
 S S R L A V H I K S L G L R A Q Q A I I P V Y F E K
 CGAAATGGGTCAATTGTTCAATGGTGGTGTCTCAAGTGTGTAATGTTTCACTTAATGATCCCAATGATCCACCA 2000
 S K W V I A S M L A V L K S G N A F T L I D P N D P P
 GCTCGAACTGCCAGTTCGTCACGAGACTGGGCGACTGTAGCGCTTACTTCCAAGCTACACCGGAGACTGTACAGAA 2080
 A R T A Q V V T Q T R A T V A L T S K L H R E T V Q K
 GCTTGTAGCGGTTCGGTGTGTTGATGACGAGCTTCTGCAATCAGTTTCTGCCAGCAGATTTCTCAAGTCTGACCA 2160
 L V G R C V V D D E L L Q S V S A S D D F S S L T
 AATCGAAGACTTGGCTAGTGATCTTCACTTCGTGACACGGCGACCCGAAAGCGATCATGATTTGAACACCGGCG 2240
 K S Q D L A V Y I F T S G S T G D P K G I M I E H R A
 TTCTCATCATGTGCACTCAAGTTCGGCGGTCTCTTGGCATCACTTGATACTCGTGCCTACAAATTGGAAACCCATGC 2320
 F S S C A L K F G A S L G I N S D T R A L Q F G T H A
 CTTTGGCGATGTCCTCTCGAGATTATGACTACTCTCATCAACGGTGGTGGCTTTGTTATTCCTCCGACGATGATCGTA 2400
 F G A C L L E I M T T L I N G G C V C I P S D D D R
 TGAACAGTATCCCGTCTTCATCAACGATCAACGTTAATTTGATGGTACGACACCTTCGTATCGGAACCTTTTCA 2480
 M N S I P S F I N R Y N V N W M M A T P S Y M G T F S

Fig. 1B

CCTGAAGACGTTCTTGCGCCTTGGCACAATTGGTACTTGTGGGGAGCAGATGTTCATCTTCACTCAAGCAATCTGGGCC 2560
 P E D V P G L A T L V L V G E Q M S S S V N A I W A P
 CAAGCTCAACTCTTCAACGGGTACGGACAGAGTGAAGTTCTTCAATTTGTTCCTTCCAATATGTCAACTGAGCCCA 2640
 K L Q L L N G Y G Q S E S I C F A S N M S T E P
 ACAACATGGGACAGACGTGGAGCTCATTCATGGGTCAATTGACCCGACGATATAAACCCAGCTAGTTCCGATGGAGCT 2720
 N N M G R A V G A H S W I D P N D I N R L V P I G A
 GTGGGAAGACTGGTCTATGAGATCCAGCATTCGCCGACTACATTTGTCCTCCCTCCGGAGAAGTCCCATTCCTT 2800
 V G E L V I E S P G I A R D Y I V P P P P E K S P F F
 CACAGACATTCGAAGTGGTATCCAGCGAACACAGTTTCTTGATGGGCAAAAACCTACAGACACAGAGATCTTCAAGAT 2880
 T D I P S W Y P A N T P F P G A K L Y R T G D L A R
 ATGCTCCGATGGGTCCATCGTTTGCCCTTGGCGCATAGACTCGCAGGTCAAGATCCGGGACACGCTGTTGAGCTGGGT 2960
 Y A S D G S I V C L G R I D S Q V K I R G Q R V E L G
 GCCATTGAGACCCATCTCCGACAGCATGCCAGCAGCTTGACTATTGTGTGAGAGCTACCAAGCGATCCCAATCTGC 3040
 A I E T H L R Q Q M P D D L T I I V V E A T K R S Q S A
 CAACACATCTCTTAATTGCAATCCCTAATAGGGTCTTCTTACTTCGAAATAGACCCCTCGGATGCCCATTTGGACC 3120
 N S T S L I A F L I G S Y F G N R P S D A H I L D
 ATGATGCTACCAAGCTATCAACATAAGCTGGAGCAGGTATTGCTCGACACTCTATCCCTCAITTCATCTGCATG 3200
 H D A T K A I N I K L E Q V L P R H S I P S F Y I C M
 CTGGAGCTTCCACGTACTGCCCGGGAAGATAGATAGGAGCGACTACGAATCATGGGCAAGACATCTTGGACAAGCA 3280
 L E L P R T A T G K I D R R R L R I M G K D I L D K Q
 GACCCAAGGGCCATTTCAACAAGCACCCGCTCCTATCCCTTCTTTCGAGACAGACAGCAAGCTCCACAGTATCT 3360
 T Q G A I V Q Q A P A P I P V F A D T A A K L H S I
 GGGTACAGAGTTTGGGTATCGATCCAGCCACGGTCAATGTGGGCAACTTTCTTGAACCTGGAGGAAACTCTATCACT 3440
 W V Q S L G I D P A T V N V G A T F F E L G N S I T
 GCTATCAAGATGGTGAACATGGGCGGTGGTATGGACCTCAAGCTCTTAACATCTACCAAGCACCCGCGCTTC 3520
 A I K M V N M A R S V G M D L K V S N I Y Q H P T L A
 GGAATTTCCGGGTGCTCAAGGTGATCTCTGTCTACACTCTATCCCAAGCAACTCATGAGGACCTGTTGAGC 3600
 G I S V A V G G D P L S Y T L I P K S T H E G P V E
 AGTCTTATTCAAGGCCCACTATGTTCTCGATCAGTTGGCAGTTGGCAGTCTGTGGTATCTGATTCATTCATGCTGTG 3680
 Q S Y S Q G R L W F L D Q L D V G S L W Y L I P Y A V

Fig. 1C

AGAATGCGGGCGTGTCAATGTCGACGCGTTACGTCGGGCTCTTGACGCGCTTGACACGGCACGAGACTCTTTAGAAC 3760
 R M G P V N D A L R A L A L E Q R H E T L R T
 GACATTTGAAGACCAAGGATGGTTCGGGTGTACAAATGTTACAGAGAGCTTTTCAGGAGATGAAGGTCATGACTCT 3840
 T F E D Q D G V G V Q I V H E K L S E E M K V I D L
 GTGGTTCAGACCTTGACCGTGTGAAACCAAGAACATCTCCCTTCAATCTCATCTCAAGAGTGGCTGG 3920
 C G S D L D P F E V L N Q E Q T P F N L S S E A G W
 AGAGCGACGGCTCTTTAGCACTTGGTGAAGATGACCACATCCTCACTATTGTTCATGCATCACTCATCTCAGATGGTTGGTC 4000
 R A T L L G L G E D D H I L T I V M H I I S D G W S
 AATTGTGTCCTTGCGACGCGATCAATCAGCTACTCAGCTGCGCTCAAGGACTCAAAAGACCCGCTGTCAAGACTCA 4080
 I D V L R D L N Q L Y S A A L K D S K D P L S A L
 CTCCTTACCTATCCAGTRACAGCACTTTGCAAAATGGCAGAAGGACCAATTCATAGAGCAGGAGAAGCAACTCAACTAC 4160
 T P L P I Q Y S D F A K W Q K D Q F I E Q E K Q L N Y
 TGAAGAAGCAACTCAAGACTCTTCCCAGCAAGATCCGACCGACTTTGCCCGCCCTGCACCTTCTCTCTGGAGACGC 4240
 W K Q L K D S P A K I P T D F A R P A L L S G D A
 AGTTTGCGTACATGTACCATCAGCGGAGCTCTACCAGTCCCTTCGAGCCTTCTGCAAGCAACACACACGACCTCTT 4320
 G C V H V T I D G E L Y Q S L R A F C N E H N T T S
 TCGTCTTCTTCTAGCTGGTTCGTCGGCTCAATTATCGTCTCAAGCTGTTGAAGACGCTGTCAATGTTGGTACACCAATT 4400
 F V V L L A F R A A H Y R L T A V E D A V I G T P I
 GCGAATCGCAACCGACTGAACCTGGAGGATATCATCGGCTGCTTTGTCAATACGAGTGTATGCGAATCAACATAGATCA 4480
 A N R N R P E L E D I I G C F V N T Q C M R I N I D H
 TCACGATACCTTTGGGACTTTGATCAACCAAGTCAAGGCTACGACGACAGCAAGCAATTCGAGACGAGGATATTTCCGTTTG 4560
 H D T F G T L I N Q V K A T T T A A F E N E D I P F
 AGCGGTGTATCAGCACTACAGCTCGATCCAGAGATCTGTCAAGCAACCTCTCGCAACACTCATTTTTCGAGTGCAC 4640
 E R V V S A L Q P G S R D L S S T P L A Q L I F A V H
 TCACAGAAGGACCTTGGAGATTCAGATTCCAGGCTCTGAGTCCGTAACCTGTGCTAGCAAGAGGTACACTCGATTGA 4720
 S K D L G R F K F G G L E S V P P S K A Y T R F D
 CATGGAGTTCATCTGTTTCAAGAAACCGACAGCGCTTAAAGTAGCGGTCAACTTTCCGATGAGCTGTTTCAAATGAGA 4800
 M E F H L F Q E T D S L K G S V N F A D E L F K M E

Fig. 1D

CTGTTGAAATGTCGTGAGAGTATCTTTGAGATTCTGAGAACGGGCTTCAAAGTTTCGGGACACCACTCTCAATCTT 4880
T V E N V V R V F F E I L R N G L Q S S R T P V S I L
CCTTTGATGATGGATCTGAGCTTGTGAAATAATGCTTCAACGTL D V L N V K H V D Y P R E S S
P L T D G I V T L L E K L D V L N V K H V D Y P R E S S
CITTTGGCTGATGCTCTCCAGACCAAGTCTCTGCTTACCCCGATAGTCTGGCTGTGGGACTCTCGTCCGATGACCT 5040
L A D V F Q T Q V S A Y P D S L A V D S S C R L T
ACACCGAGTTGGATCGCCAGTCTGATATCTCGCTGATGCTGACGCTCAATGCTGCAGAGACGCTTGTTCGA 5120
Y T E L D R Q S D I L A G W L R R R S M P A E T L V A
GTATTTGCCCCACGGTCACTGTGAGACAAATGTCGCTTCTTTGGTGTGAGGGCAACTTGGCTATCTTCTCTCGA 5200
V F A P R S C E T I V A F F G V L K A N L A Y L P L D
TGTACGATCGCCCTCGGCGAGATTCAGGATATACCTTCTGGACTTCTGGGCTTACCATTTTGTGTTGTTGATGATGATA 5280
V R S P S A R V Q D I L S G L S G P T I V L I G H D
CAGGCGCTCCCGATATCGAGGTTACTAACGTCGAGTTTGTTCGTATCCGGGATGCGCTGAATGACAGCAATGCGATGCG 5360
T A P P P D I E V T N V E F V R I R D A L N D S N A D G
TTTGAAGTCATCGAGCACACAGCAAAAGCCCTCAGCCACGAGTCTCGCATACGCTGTATACCTCAGGATCCACTGG 5440
F E V I E H D S T K P S A T S L A Y V L Y T S G S T G
CCGACCAAAAGGCGTATGATTGAGCACCGTGTCATTTTCGAACAGTCACAAAGTGCGCTGTATACCAACTATCTCTCGG 5520
R P K G V M I E H R V I I R T V T S G C I P N Y P S
AAACGAGGATGGCTCACATGGCGACCATTCGTTTCAACGGCGCATCGTACGAGATCTACACGCGCCCTTTTGTTCGGAAGG 5600
E T R M A H M A T I A F D G A S Y E I Y S A L L F G R
ACACTTGTTCGTTGACTACATGACAAACCCCTCGACGTAGACCACTCAAGATGTGTTTTCCGAGAGCATGTCAACGC 5680
T L V C V D Y M T T L D A R A L K D V F F R E H V N A
GGCAAGTCATGTACCAGCTCTTCTCAAGATGATACCTCCGAGTCCGAGAGGCTTCGAGAACCTTGATCTCTTCT 5760
A S H V T S S Q D V P L R V P R L S R T L M F
TCTTTGGTGTGACAGATTCGACGGCCCCAGATGCTTCGATGCGCAGGCTTTATCAAGGGTCCAGTGTTCACATGGT 5840
F L V V T D S T A P D A Q G L Y G V Q C Y N G
TACGGCCCAACAGAAATGGAGTCATGAGTACAAATCTATCCCATGTAGTCGATCGATCAATGAGGAGTCCCAAT 5920
Y G P T E N G V M S T I Y P I D S T E S F I N G V P I

Fig. 1E

TGACGAGCTCTGAACAACACTCAGGAGCGTATGTCGTGGATCTGAGCAACAGCTTGTGGCATTTGGTGTGATGGAGAGC 6000
 G R A L N N S G A Y V V D P E Q Q L V G I G V M G E
 TTGTTGACATGGCGATGFTCTTGGCGGGTACAGTGACAAACCCCTTTGTCGACATTAATCTGTC 6080
 L V T G D G L A R G Y S D K A L D E N R F V H I T V
 AATGACCAGACAGTGAAGCGTATCGCACTGGCGATCGAGTCCGGTACAGGATGGAGATGGCCCTCATCGAGTCTTTCGG 6160
 N D Q T V K A Y R T G D R V R I G D G L I E F F G
 ACGTATGGACACCCAGTTCAAAGATCGTGGCAATCGATCGAATCAGCTGAGATGAAGCGGCCCTTCTCGCGCACTCCT 6240
 R M D T Q F K I R G N R I E S A E I E A L L R D S
 CCGTCCGAGATGCTGCTGCTCTTCCAGCAAGATGAGGATCAAGCGCTGAGATCTTGGGTTTCTGTTGCTGATCAT 6320
 S V R D A A V L Q Q N E D Q A P E I L G F V V A D H
 GATCATTTGAGAATCACAGGGACAATCTGCCAATCAAGTGAAGGATGGCAAGACCAATTCGAGAGTGGCATGTATTC 6400
 D H S E N D K G Q S A N Q V E G W Q D H F E S G M Y S
 CGACATTTGGCGAAATTGACCCGTCGACGATTTGGTAGCGACTTCAAGGGTTGGACATCAATGTATGATGGAAGTCAATCG 6480
 D I G E E I D P S T I G S D F K G W T S M Y D G S Q I
 ACTTCGATGAGATGACGAGTGGCTTGGTGAGACTACCCGACACTCCATGACAAATCGCTCTTAGCAATGTCTTGAA 6560
 D F D E M H E W L G E T T R T L H D N N R S L G N V L E
 ATTGGAACAGGTAGCGGATGATCCTCTTCAACCTTGACAGCAGGCTTGAGAGTTAGCTTGGTCTTGAACCATCCAGATC 6640
 I G T G S G M I L F N L D S R L E S V V G L E P S R S
 AGCAGCTGATTTGTCAACAAAGCTACGAGTCTATACCATCGTTGCTGGAAAGCCCAAGGTTTCAGGTTGGAAACAGCTA 6720
 A A A F V N K A T E S I P S L A G K A K V Q V G T A
 CAGATATTGFTCAAGTCGATGACTTACACCCGTGACCTCGTGGTTCTCAACTCAGTCATTCAGTATTCCTCGTCTTCGGAG 6800
 T D I G Q V D D L H P D L V L N S V I Q Y F P S S E
 TACTTTCAGAAATCGACACACTTGTATTCATCTGCCTAACGTGACGGATTTTCTTTGGCGATGTCGGATCGCAGGC 6880
 Y L E I A D T L I H L P N V Q R I F F G D V R S Q A
 CACCAACGAGCACTTCTTGTCTCCAGGCTTATCCACACTGGGGAGAAATGCAACAGGACGATGTTTCGACAGAAA 6960
 T N E H F L A A R A I H T L G K N A T K D D V R Q K
 TGGCAGAATTTCGAGACATGGAGGAGGATGCTTGTGTAACCTGCTTTCACCTGTTGAAAGACAGGTTTCCAGGT 7040
 M A E L E D M E E L L V E P A F E T S L K D R F P G
 CTCGTGGAACATGTTGAGATCTCGCCAAAGAACATGGAAGCTGTGAATGAGCTCAGTGGCTATGCATGCGCGCTGTGT 7120
 L V E H V E I L P K N M E A V N E L S A Y R Y A A V V

Fig. 1F

GCACGTTCCGGGTTCACTTGGAGATGAGCTTGTGCTTCCGGTTTGAGAAAGATGACTGGATCGACTTTCAAGCGAATCAAT 7200
H V R G S L G D E L V L P V E K D D W I D F Q A N Q
TGAACCAAGTCACTGGGTGACCTTCAAGTCTTCAGATGCTGCTATCATGCGAGTCAGCAAAATCTTTTCGAAATC 7280
L N Q K S L G D L L K S D A T A I M A V S K I P F E I
ACGGCTTTGAAAGACAGTTCGTCCCTCAATAGCAACATCGATGAGTGGCAGCTATCAACCATTCGGTCCAGCGC 7360
T A F E A R Q V V A S L N S I D E W Q L S T I R S S A
CGAGGCACTCATCATCTCCGTTCCGACACATCTTCGCATTGGGGAAGCCGGTTCCGTGTCGAGTCACTTCTG 7440
E G D S S L S V P D I F R I A G E A G F R V E V S S
CACGACATGGTCTCAGAATGGTGCATTGGACGCTGTTTCCATCATGTTGTCTCCAAGGCGTACTCTGGTCAACTTT 7520
A R Q W S Q N G A L D A V F H C C S Q G R T L V N F
CCTACGGACCATCACCTTCGAGGGTCTGATCTCTCACCATCGACCCCTTCAGCGACTGCAAAACCGTGTATCGCCAT 7600
P T D H H L R G S D L L T N R P L Q R L Q N R R I A I
CGAAGTCCGCGAGAGGCTTCGGTCCCTTACTTCATCTGATATCCCATCGAACATCGTTCTCTGGACAAGATGCCTC 7680
E V R E R L R S L L P S Y M I P S N I V L D K M P
TCAAGCCCAATGGTAAAGTTGACCGGAAGAACTCTCTCGCAGGGCAAGGTTGTACCGAAGCAGCAGCAGCGCGG 7760
L N A N G K V D R K E L S R R A K V V P K Q Q T A A P
TTACCGACATTTCCCATCAGTGAAGTCAATCTTTTCGGAAGAAGCCACTGAGGTGTTTGGCATGAAGTTGACAT 7840
L P T F P I S E V E V I L C E E A T E V F G M K V D I
TACCGATCACTTCTTCAATCTCGGTGGACACTCTCTCTTGGCCACGAGCTCATTTCTCGTATCGACCAACGACTCAAGG 7920
T D H F F N L G G H S L L A T K L I S R I D Q R L K
TCCGTATCACTGTCAAGGATGCTTTTGACCATCTCTGTATTTGGGAGTATGACATCTGTATCCGTCAAGGCTGGGTTTG 8000
V R I T V K D V F D H P V F A D L A S V I R Q G L G L
CAACAACCCGTTTCTGATGGTCAAGGCAAGACAGATCTGCCCATATGGCACCCCGTACCGAGACTGAAGCTACTCTG 8080
Q Q P V S D G Q G Q D R S A H A M A P R T E A I L C
TGATGATTTGCAAGGTTCTGGGGTTCTCAAGTCCGGGATACAGAAATTTCTTGATCTTGGTGTCACTCACTCAAG 8160
D E F A K V L G F Q V G I T D N F F D L G H S L M
CTACTAAACTCGCTGTGGCGCATCGACATCGACTTGACACAGCACTGTTTCGGTGAAGGATGTTTTCGATCATCTCTG 8240
A T K L R I G H R L D T T V S V K D V F D H P V L
TTCCAACCTTGCATTTGCATTGGATACTTGGTTCAATCCAGACCAATGAGATAGTTGGAGGTAGAGAAATGGCTGAATA 8320
F Q L A I A L D N L V Q S K T N E I V G G R E M A E Y

Fig. 1G

CTCACCTTTCCAACCTCCTCTTTACAGAAGACCAGAGGAGTTTATGGCGAGCGAGATCAAGCCACAACTTGAGTTACAGG 8400
 S P F Q L L F T E D P E E F M A S E I K P Q L E L Q
 AAATCATTTCAAGACATATATCCGCTTACCAGATGAGAGGCTTTCTCTTCGATCACAACTGCGCGCCGACCT 8480
 E I I Q D I Y P S T Q M Q K A F L F D H T T A R P R P
 TTCCTGCGGTTCTACATCGACTTCCCGACACTCCAGCCTGATGCTCGAGGTCTAATCAAGGCTTGCGAGTCTCTGGT 8560
 F V P F Y I D F P S T G S E P D A A G L I K A C E S L V
 AAATCATCTTGACATCTTCAGAACAGTCTTTGCAGAGGCATCTCGAGAATATACCAAGTGGTCTTGCTCTGTCTTGATC 8640
 N H L D I F R T V F A E A S G E L Y Q V L S C L D
 TGCAATCCAAGTGATTGACACAGAAGACAAACATCAATACGGCGCAAAATGAGTTTCTCGATGAGTTTGGCAAAGCCA 8720
 L P I Q I E T E D N I N T A T N E F L D E F A K E P
 GTTCGCTGGGACATCCGTTGATTGCTTTACAATCATCAAAACCAAGTCGATGCTGTGATAATAGAAATATCGCA 8800
 V R L G H P L I R F T I K Q T K S M R V I M R I S H
 TGCCCTGTATGATGGTCTGAGTCTAGAGCATGTCGTGCGCAAACTTCACATGCTCTACAACGGGAGATCACTTTTGGCAC 8880
 A L Y D G L S L E H V V R K L H M L Y N G R S L L P
 CACACCAATCTCGCGTACATGCAGTATACCTGACGGTCGCGAAAGTGGACATGGATTTTGGCGGATGTGATTCAA 8960
 P H Q F S R Y M Q Y T A D G R E S S G H G F W R D V I Q
 AATACGCCCATGACAAATATTGAGTGATGACACGCTTTGATGGAATGATGCAACCTGCAAGGCTTGCACCTATCAA 9040
 N T P M T I L S D T V D G N D A T C K A L H L S K
 GATTTGCAATATTCCTTCACAGGTACTTCGAGGCAGCAGTAAACATCACTACTCAAGCTACTGTGTGTTAACGCGACCTGG 9120
 I V N I P S Q V L R G S N I I T Q A T V F N A C
 CGTTAGTCTTGTACGCGGATCTGACTCGAAAGACGTTGCTTTGGACGATCGTCTGTGTCGCAAGCTTCGCTGTT 9200
 A L V L S R E S D S K D V V F G R I V S G R Q G L P V
 GAATACAGGACATTTGCGGCTTTGTACCAACGCGATTCCTGTTGCGCTCATATAGAGTCGTCAGATTACACCAATT 9280
 E Y Q D I V G P C T N A V P V R A H I E S S D Y N Q L
 GCTGCGACATCCAAAGACAGTACCTTCTCAGCTTCCACACCAACAAATTGCTCTCAGATCTCAAGCNACTGTA 9360
 L H D I Q D Q Y L S L P H E T I G F S D L K R N C
 CAGATTGCCAGAAGCAATCACCAACTTCTCATGCTGCATCAACATCCACAAATTCAGTACCACCCGAGAGTCAGTTC 9440
 T D W P E A I T N F S C C I T Y H N F E Y H P E S Q F
 GAACAGCAGAGGTTGAGATGGTGATATGACAAAGTTTGTCAACATGAGATGGAGCCATATATGATTGGCCAT 9520
 E Q Q R V E M G V L T K F V N I E M D E P L Y D L A I

Fig. 1H

TCGCGGTGAAGTTGAACACAGACGGAGCAGGACTGAAGGTTTACTGTTATCGCGAAGACGACAGTATTTGGTAGGAAGACAG 9600
 A G E V E P D G A G L K V T V I A K T Q L F G R K R
 TAGAACATCTGTGTGGAGGAAGTTTCCAAAACGTTTGAGGGTCTCAACTCTCTTTGTAACGCACGCGGTTGGTCTCAATCG 9680
 V E H L L E E V S K T F E G L N S S L . . .
 TCGCGACAGAACACCGATAGGTTTGTAACTTCTTAATGACGCTTTGACTTTTGGTTTTCATTCGGAGCAATA 9760
 GTAAGCAACACTCCGAAATGCAGATATTACACTTCAGAACTATTACTTGACTATTATCTACGTTGTCAAGTTTCA 9840
 CATGCTTCTAGCTTCGATCGAGTCAAAAATTGAGATCTACAGGTTAACGCGAAATCCAGAACAAATTCACAGGATTCA 9920
 TCGATCGAACACTATGATTGGTTCGGTCTTCGACAGGACCATTTGTCCAAATAGAAGATAGATAAGATATGCGAG 10000
 AATCGCAAGGATTGGAACATCCGATCAGCTTGATCTAGTCCCTAAACAGTAAATAGTCTGGCTTCTTCGGATTTCG 10080
 CTTTGGTGTATGAATTATCATAGTTTGTTCGAAAGAACGATGACAGCTTGTATTTTGGATATATATAAGTTTC 10160
 ATAAAGGTATGACTCTTGATGATGAATCAATTAGAACAATACCTTGAACAATATGTGTCCAACTATACAACTTGA 10240
 ACGGACACTCTTCAACATCAACAATGGATCTGTCCAAAGAAGCTGCCTACCTGTTCAGTCCAAATGGGCCACCAT 10320
 CGAAGTCAGATCTGCACCTGTTCACAACCTGGTTCAGGAGAGTTGCTTATCAAGACACATGCTGTCCGATCAATCCAG 10400
 TCGACGGAGTGAAGAGTCCATGGGTAAATGATGTTGAATGGCTCAAGTATCTCTCATCTTTGGCTACGATGTCCGT 10480
 GCGAGGTCATCAAGACGGGACCTTGGTGTAGTTCGATTTAAAGAAGGGGATAGATGTGGTGCTACAGCAGCATGGA 10560
 CAAGCGAAGAAGATCCGACGAAGGCGCATTTCAAGAAGTTTGCAATCATCGAGAGCATTTGGCTGCTCGAATTCAG 10640
 AGCGTTTACGTCACCGATCCGACGGTTTTCGCTCTGACTTCGTACGGGTGCACTGTGCTTCCAAAAGGATCAA 10720
 CTGGCATACAACTACTCAAAACCAAGTCAAGCGCAGTGCACAAGTCAGACATGTGTTTGGGAGCGGATACAA 10800
 TGTTCGGAGAATGCTGACAGCTTGTCTCGCGCCGGCTATGATGTTGTCCGACAGCATCACCTAAGAACTGGGATA 10880
 TCGTACGGGTCTCGCGCTTGTGCAAGTTTGTGACTATCAAGTCTATCGGCCATAAAGCATGTGGTATATCTGCTTTCAAG 10960
 GACAAGAAATGCGAGGTGCTGTAGCTATTTGGTCAAGGGTCACTTGGCGAAATCGTCGACATGTCAAGCGTTCGGG 11040
 AGCCACCAAGAATGTTGCGCAAGTTACCTCTCAATGCGCTGAGTTCAGACCAACACCAAGATATCCATGATTCCTGTTG 11120
 TCGCAAGATATTTCTGGATGCGCGGAACATGATCGACTCAAGGTTTCGAGCTGGAGTCCAAAGCAAGTTTGTTTTGGT 11200
 ACAGACATAATT 11212

Fig. 11




Fig. 2

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Methods For Producing Polypeptides In Cyclohexadepsipeptide-Deficient Cells

the specification of which (check only one item below):

☐ is attached hereto

☒ was filed as United States application

Application No. to be assigned

on January 13, 2000

and was amended

on _____

☐ was filed as PCT international application

Number _____

on _____

and was amended under PCT Article 19

on _____

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim priority benefits under Title 35, United States Code, §119 of any provisional or foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR U.S. PROVISIONAL/FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY	DATE OF FILING	PRIORITY CLAIMED
		<input type="checkbox"/> YES <input type="checkbox"/> NO
		<input type="checkbox"/> YES <input type="checkbox"/> NO
		<input type="checkbox"/> YES <input type="checkbox"/> NO

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY
(Includes Reference to PCT International Applications)

Attorney's Docket Number:
5778.200-US

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this applications is not disclosed in that/those prior application(s) as the matter provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:

U.S. APPLICATIONS		STATUS (Check one)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	Patented	Pending	Abandoned
09/229,862	January 13, 2000		X	
PCT APPLICATIONS DESIGNATING THE U.S.				
APPLICATION NO.	FILING DATE	US SERIAL NUMBERS ASSIGNED (if any)		

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Steve T. Zelson Reg. No. 36,335 Elias J. Lambiris Reg. No. 33,728 Valeta A. Gregg Reg. No. 35,127 Carol E. Rosek Reg. No. 36,993 Robert L. Starnes Reg. No. 41,324 Resa Green Reg. No. 38,475

Send Correspondence to: Steve T. Zelson, Esq.
Novo Nordisk of North America, Inc.
405 Lexington Avenue, Suite 6400
New York, New York 10174-6400

Direct Telephone Calls To:
Steve T. Zelson
(212) 867-0123

1	Full Name of Inventor	Family Name Berka	First Given Name Randy	Second Given Name M.
	Residence & Citizenship	City Davis	State or Foreign Country California	Country of Citizenship United States
	Post Office Address	Post Office Address 3609 Modoc	City Davis	State & Zip Code/Country California 95616/US
2	Full Name of Inventor	Family Name Rey	First Given Name Michael	Second Given Name W.
	Residence & Citizenship	City Davis	State or Foreign Country California	Country of Citizenship United States
	Post Office Address	Post Office Address 605 Robin Place	City Davis	State & Zip Code/Country California 95616/US
3	Full Name of Inventor	Family Name Yoder	First Given Name Wendy	Second Given Name T.
	Residence & Citizenship	City Winters	State or Foreign Country California	Country of Citizenship Great Britain
	Post Office Address	Post Office Address 8503 Pleasants Valley Rd.	City Winters	State & Zip Code/Country California 95694/US

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY
(Includes Reference to PCT International Applications)

Attorney's Docket Number
5778.200-US

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1



Date

June 11, 2000

Signature of Inventor 2



Date

1/11/00

Signature of Inventor 3



Date

1/11/00

09492798-011300

SEQUENCE LISTING

<110> Randy M. Berka
Michael W. Rey
Wendy T. Yoder

<120> Methods For Producing Polypeptides In
Cyclohexadepsipeptide-Deficient Cells

<130> 5778.200-US

<140> To Be Assigned

<141> 2000-01-13

<150> 09/229,862

<151> 1999-01-13

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 11212

<212> DNA

<213> Fusarium

<400> 1

aattagattc	cactagtacg	ccattgtaga	atcaaggcca	agatatgaac	aaccataag	60
taacggcgat	cctgtctcat	gtatccaaaa	ataagagaca	cggcataatc	actgctttgc	120
agatctttct	tcaaatctct	ccctcgagaa	gctactggga	tgaatgagtc	tcttggctca	180
gattagatat	attcaactgta	tctgcgaat	agactttgcc	tggtagcatt	aacgttccta	240
tattctatta	tcaaatctct	acattcaata	tggaaatctc	tactgctgtc	gatggtaggc	300
aagacctgcc	acctacacca	gcttctgttt	gtagtcatgg	agatagtcct	ctcaatagct	360
cttacgagca	actcttccat	ctctatggtc	tggattcgag	tcgcatacga	gctatcaaac	420
catgcacacc	tttccagctt	gacatgatgc	actgcaatgc	tttgataatc	cagtcctgcta	480
tccggccatgc	gggtgatgat	gtcccaaccg	acattgacat	ctctcgtttc	gcgcttgctg	540
ggaaggagat	cgtaaccaca	acccagacct	tgcgagcctt	tgccttcacc	tcggactctg	600
gaaagacttc	tcaagtcatc	ctaaaaagata	gctttgtctt	ctcatggatg	tgttggtctt	660
cttcgagctc	ccagatgaa	gtgggttcggg	atgaagctgc	cgctgctgca	tccgggccac	720
gctgcaacgc	ctctgttcta	cttgaagaca	tgcagacgaa	gaaatgtcag	ctgggtttgga	780
cttctcagca	tgcattggta	gacgtcaactt	tccaacaacg	cgctcctgagc	cggtgtttcg	840
cggtctacaa	gcatagagag	gacacacatc	ggcctgagac	acccagagtc	tctgatgcca	900
ctgacactga	ctctcagtc	gtctccgtgg	gtctcatgag	ctgcgaggac	aatgcctgat	960
cggtcgactca	ttctctgcaa	actcaacctta	acgatctcaa	tgcgtccgtc	tccctccacc	1020
tgtcttgacca	cctgatgggt	cccaaccaca	ctacaacagc	agagcatcgt	atcacattcc	1080
ctctttcaca	gaaagcacta	tccaattctg	ccatctgcgc	tactgcactc	tcaatactcc	1140
tctcgcgcta	caactcaact	gacgaggcct	tgtttggtgc	ggtaactgag	caatctctac	1200
catttgacaa	acactatctt	gcagatggta	cgtaaccaac	agttgacacc	cttcgtgtac	1260
actgccaatc	aaatctctgt	gcatacagat	tcatggatgc	aatctctctt	tacgatgatc	1320
gccttggtca	ctctgcacca	tttggccttc	gcgacatccg	caacactggg	gataatggct	1380
ctgcgcctgc	cgatttccaa	actgttttac	tctgcaccga	tggcagccac	gtaaaccaatg	1440
gtatccacgg	tttccctcaa	cagataaacag	agtcacagcca	ttctatgcct	tgaacacacc	1500
gtgcctctct	tctgcaactg	cagatggaaa	gtacgggagc	ctctgctgggt	gcctactatg	1560
accacaaatg	tatcgattgt	ctctcagacaa	cgctctgctg	acagcagttt	gtctactctga	1620
tcaagtgttt	gcaaagtcct	ctagacctga	gctctatggc	tgaggtcaac	ttgatgactg	1680
agtatgacag	agcagagatt	gagagttgga	actcgcaacc	gttagaggtg	caggataacc	1740

tgatccacca	tgagatgttg	aaagctgttt	ctcattcccc	cacccaaaacg	gccatccaag	1800
cgtagggatg	agactggacc	tattccggagc	tgcacaatgt	ttcgtaacga	ctcgctgtcc	1860
atatacaagt	actttggcctt	agagctcagc	aaagccattat	tcagatctct	tttgagaagt	1920
cgaatagggt	cattgtcttca	atgtctggctg	ttctcaagtc	tggtaatgct	ttcaactctaa	1980
ttgatcccaa	tgatccacca	gcgcgaactg	cccaggtcgt	ccagcgactc	cgggcgactg	2040
tagcgcttac	ttccaaagcta	caccgcgaga	ctgtacagaa	gcttcttaggc	cgttgcgttg	2100
tggttgatga	cgagcttctg	caatcagttt	ctgcacagca	cgattttctc	agctcgacca	2160
aatcgcaaga	cttggcctac	gtgatcttca	cttctggtag	cacggggcag	ccgaaaggca	2220
tcatgattga	acacgcagcg	ttctcatcat	gtgcactcaa	gttcggcgcg	ttctttggca	2280
tcactctcga	tactcgtgcc	ctacaatttg	gaacccatgc	ctttggcgca	tgtcttctcg	2340
agattatgac	tactctcact	aacgggtggct	gcgtttgtat	tcctctccag	gatgatcgta	2400
tgaacagatg	ccgtctcttc	atcaaccgat	acaaagcttaa	ttggatgatg	ggagacacct	2460
cgtacatggg	aacctttttca	cctgaagacg	ttcttggtct	tgccgacattg	gtactttgtt	2520
gggagcagat	gtcatcttca	gtcaacgcaa	tctgggcccc	caagctccaa	ctcttgaacg	2580
ggtagcgaca	gagtgaaagt	tcttcaattt	gttttgcctc	caatatgtca	actgagccca	2640
acaacatggg	cagagcagtc	ggagctcatt	catgggtcat	tgacccgaac	gatataaac	2700
gactagtctc	gattggagct	gtgggagaa	tggtcattga	gagtcacagg	attgcccgcg	2760
atcacattgt	ttccccctct	coggagaagt	ccccattctt	cacagacctt	ctcttggcgt	2820
actacgcgaa	caagctttctc	gatggggcaa	aactctacag	gacaggagat	cttgcaagat	2880
gagcctccga	tggtgtccat	gttttgcttg	ggcgcataga	ctcgaggtc	aaagctccgg	2940
tgacagctgt	tgagctgggt	gccatttgaga	cccatctccg	acagcagatg	ccagacacct	3000
tgactattgt	ggtagaagct	accaaagcat	cccaatctcg	caacacgaca	ttcttaattg	3060
catctctcgt	aggggtctct	tacttcggaa	atagacctc	ggatgcacca	ctcttggacc	3120
atgatgctac	caaaagctatc	aaactaaagc	tgtagcaggt	attgcctcga	caactctatc	3180
ctctcattct	catctgcctc	ctggagcttc	caactactgc	caccgggaag	atagatagga	3240
ggcgactacg	aatcatgggc	aaagacatct	tggaacaagc	gaccaagagg	gccattgttc	3300
aaaacagcacc	cgctctctat	ctcgttttct	cagacacagc	acaaaagctc	ccagatattc	3360
gggtacagag	tttgggtatc	gatccagcca	cggtcaatgt	tggggcaact	ttcttcgaac	3420
tcggagggaaa	ctctatcact	gctatcaaga	tggtgaacat	ggcgagggtc	tttgggtatg	3480
acctaagggt	ctctaaacat	taccagcacc	cgacagcttc	gggaatttcc	ggggtgtcgg	3540
aggggtgacc	ttctgtctac	actctcatcc	ccaagtcaac	tcattgagga	actgtgtgagc	3600
agttctattc	acaaaggcga	ctatgggttc	tggaatcagt	ggacgtttgc	cgtctgtgtg	3660
atctgtattc	atatgctgtg	agaatgcgcg	ggcctgtcaa	tgctcgacgcg	ttaactcggg	3720
ctcttgcagc	ctgtgaacga	cgacacgaga	ctcttagaac	gacatttgaa	gaccaggatg	3780
gtgtcgttgt	acaaatttgt	cacgagaagc	ttcttgagga	gatgaaggtc	attgatctct	3840
ttggctcaga	ctctgaccgc	tttgaggtgt	tgaaccaaga	acagactact	cccttcaatc	3900
gtctcatctga	agctggctgg	agagcgacgc	tcttaacgact	tggtgaaagt	gaccacatcc	3960
tcactattgt	catgcatcac	atcatctcag	atggttggtc	aattgtatgt	ttgcgcacgc	4020
atctcaatca	gctctactca	gctgcgctca	aggactcaaa	agaccgcgtg	tcagcactca	4080
ctctctacc	tatccagtat	agcgactttg	caaaatggca	gaaggagcaa	ttcatagagc	4140
aggagaagca	actcaactac	tgggaagaagc	acctaacaaga	ctcttcccga	gcgaagatcc	4200
cgacagcagc	tgccccctct	gcaacttctgt	ctggagacgc	aggttgcgta	catgttacca	4260
tcgagggctga	gctctaccag	tccttctcag	ctctctgcga	cgaacacaca	accactctct	4320
ctgtcgcttc	tctagctcgc	ttcgttgcgc	ctcattatcg	tctcacagct	gttggaagact	4380
tctgatattg	tacaccaatt	gggaatcgca	accgactctg	actggaggat	atcatcgctc	4440
gctttgtcaa	tacgcagtgt	atgcgaatca	acatagatca	tcacgatacc	tttgggactt	4500
tgatcaacca	agtcgaagct	acgacgacag	cagcattcga	gaacgagatg	attccgtttg	4560
agcgcggtgt	atcagcacta	agcctggat	ccagagatct	gtcaagcaca	cctctcgacc	4620
aaactcattt	tgcaagtgc	tcacagaagg	accttggaag	attcaagttc	cagggtctcg	4680
agtcctgata	tgtgctcagc	aaagcgtaca	ctcgatttga	catggagtgc	catctgttct	4740
aaagtacgaa	cagcctctaaa	ggtagcgtca	acttctgcga	tgagctgttc	aaaatggaga	4800
cgttttgaaa	tgtcgtcaga	gtattctctg	agattctcag	aaacgggctc	caaatgtcgc	4860
ggacaccaagt	ctcaatcatt	cctttgactg	atggcattgt	gaactctgaa	aaattggatg	4920
ttctcacaagt	caaacatgtc	gactatcccc	gagaatcgag	cttggctgat	gtcttcacga	4980
ccccagttct	tgtctacccc	gatagtctgg	ctgtggtgga	ctctctgtgc	cgattgacct	5040
acacgcgctc	ggatcgccag	ctgatattc	tcgctggatg	gcttcgtcga	cggctcaatg	5100
ctgcagagac	gcttgcgcga	gtatttgccc	caaggtcatg	tgagacaatt	gtcgcgttct	5160
ttggtgtgtt	gaaggcgagc	ttggcctatc	ttctctcga	tgtagcatcg	ccttcggcga	5220

gagttcagga	tatactttct	ggactttctg	ggcctaccat	tgttttgatt	ggccatgata	5280
cagcgctccc	cgatattcgag	gttactaacg	tcgagtttgt	tcgtatccgg	gatggcgctga	5340
atgacagcaa	tgcagatggc	tttgaagtca	tcgagcacga	cagcacaaa	ccctcagcca	5400
cgagttctgc	atacgtgctg	tatacctcag	gatccactgg	ccgaccacaaa	ggcgctcatga	5460
ttgagcacga	tgctattatt	cgaacagtca	caagtggctg	tatacccaac	tatccttcgg	5520
aaacaggagat	ggctccacatg	cgacaccattg	cgtttgacgg	cgcatctgtac	gagatctaca	5580
gcgcctctttt	gtctggaaagg	acactttgtt	gcgttgacta	catgacaacc	ctcgacgcta	5640
gagcactcaa	ggatgtgttt	ttccgagagc	atgtcaacgc	ggcaagtcac	gtcaccagct	5700
ctctccaaag	tgtacctctc	cgagtcocga	gaaggctctc	gagaactctt	atgttctctt	5760
tgtctgttgt	gacagatttcg	atgcgtctcg	atgcgtctcg	tgcgacggga	ctttatcaag	5820
gggtccagtg	ttacaaatgtt	tacggcccaa	cagagaaatg	agtcactgagt	acaaactctac	5880
ccattgactc	gactgagctg	ttcatcaatg	tggacagact	tggacagact	gtcaacaact	5940
caggagcgta	tgtcgtggat	cttgagcaac	agcttgtttg	cattgtgtgtg	atgggagagc	6000
ttgttgctac	tggcgatggt	cttgccgggg	gctacagtga	caaaagccctt	gacgagaacc	6060
gttttgcga	cattactgtc	aatgaccaga	cagtgaaggc	gtatcgcaact	ggcgatcgag	6120
tgcggtacag	gattggagat	ggcctcatcg	agttctctcg	acgtatggac	accagttcca	6180
agattcgtgt	caatcgtatc	gaatcagctg	agattgaagc	ggcctctctg	cgagactcct	6240
ccgtccgaga	tgtctgtctg	gtccttcagc	agaatgagga	tcgaagccgt	gcgtctctgg	6300
gtgtgttgt	tgtctgattc	gatcattctg	agaatgacaa	gggacaactc	gccaatcaag	6360
tgcgaagatg	gcaagaccat	ttcgagatgt	gcactgtatc	cgacatctgg	gaaattgacc	6420
ctgcagcatg	tggtagcgac	ttcaagggtt	ggacatcaat	gtatgtagga	agtcacaaatg	6480
actctgatga	gatgcacag	tggcttggtg	agactaccgc	gacactccat	gacaaatcgct	6540
ctctagggcaa	tgtccttgaa	attggaacag	gtacggcgat	gactcctctc	aaccttgaca	6600
cgaggcttga	gagttacgtt	ggtcttgaac	catccagatc	agcagctgca	tttgtcaaca	6660
aaagtcacga	gtctatacca	tcgcttgctg	gaaaagccaa	gggttcaggtt	ggaaacagta	6720
cagatattgg	tcaagtctgat	gacttacacc	ctgacctcgt	gggtctcaac	tcagtctctc	6780
agtaattccc	gtctctcgag	taccttgcag	aaatcgacga	catctgctca	catctgccta	6840
acgtgcagcg	gattttctctt	ggcgatgtcc	gatcgcaggc	caccacacag	cacttctctg	6900
ctgcaggggc	tatccacaca	ctgggggaaga	atgcaacgaa	ggacgatgtt	tcgacagaaa	6960
tggccaatct	ggaggacatg	gaggaggagt	tgtcttgtag	actgtcttc	ctcacctcgt	7020
tgaagacag	gtttcccgagt	ctgggtggaac	atgttgagat	cctgcacaa	aaactgggaag	7080
ctgtgaatga	gtccagtgct	tatcgatatg	ccgctgttgt	gcacgttcgg	ggttcacttg	7140
gagatgagct	tgtgcttcog	gttgagaaag	atgactggat	cgactttcaa	gcgaatcaat	7200
tgaaacacaa	gtccactgggt	gacctctcga	agtcctcaga	tgtcgtctca	atggcgatca	7260
gcaaaattcc	tttcgaaatc	acggcctttg	aaagacaggt	cgctcgcttc	ctcaatagca	7320
acatcgatga	gtggcagcta	tcaaccattc	ggctccagcgc	cgagggcgcg	tcatcactat	7380
ccgttccgga	ctactcttcg	attgctgggg	aagccgggtt	ccgtgtcgag	gtcagttctg	7440
cacgacagtg	gtctcagaat	gggtgcattg	acgtgtttt	ccatcatgtg	tgtcccaag	7500
ggcgtaactc	gttcaacttt	cctaaggacc	atcaccctcg	aggggtctgat	ctcctcaaca	7560
actcgacccct	tcacgagctg	caaaaaccgc	gtatgcocat	cgaagtccgc	gagaggcttc	7620
ggctcttact	tcacatcgac	atgatcccat	cgaacatcgt	tgttctggag	aaagatgcctc	7680
tcaacgccaa	tggtaaagt	gaccgggaag	aaactctctg	caggggcaag	gttgtaacga	7740
agcagacac	agcagccgcg	ttaaccgacat	ttcccatcag	tgaagtcgaa	gtcatctctt	7800
gcgaagaagc	cactgagctg	tttgcatga	aggttgacat	taccgatcat	ttcttcaatc	7860
tcgggtgaca	ctctctcttg	gccacgaagc	tcattttctg	tatcgaccaa	cgactcaagg	7920
tcggtatcac	tgtcaaaagt	gtctttgacc	atcctgtatt	tgcggatcta	gcactctgca	7980
tcgctcaagg	gctgggttgt	caacaaccgc	ttctctatgg	tcagggaacaa	gacagatctg	8040
ccacatggc	accocgtacc	gagaactga	ctatactctg	tgtatgagtt	gcaaaaggtt	8100
tggggttcca	agtcgggatt	acagacaatt	ttcttgcatt	tgggtgtcat	tcactcatcg	8160
ctactaaact	cgtctgtcgc	atcggacatc	gacttgacac	gactgtctcg	gtgaaggatg	8220
ttttcgatca	ttctgtactc	ttccaacttg	caattgcatt	ggataactctg	ttctcaatca	8280
agaccaatga	gatagttgga	ggtagagaaa	tggctgaata	ctcaccttct	caactcctct	8340
ttacagaaga	cccgaggag	tttatggcga	cgagatcaaa	gccacaactt	gagttacagg	8400
aaactattca	agacatatat	ccgtctaccc	agatgcagaa	ggctttctct	ttcgatcaca	8460
caactgcgcg	cccgagacct	ttcgtccggt	ttctacatga	cttcccagcg	acttccgagc	8520
ctgatctgcg	aggtcttaact	aaagcttgcc	agtcctctgg	aaatcatctt	gacatcttca	8580
gaacagctct	tgcagaggca	ctctggagaa	tataccaagt	ggctctgtcc	tgtcttgatc	8640
tgcacatcca	agtgatgtgag	acagaagaca	acatcaatac	ggcgacaaat	gagtttctcg	8700

atgagtttgc	gaaaagagcca	gttcgtctcgg	gacatccggt	gattcggtttt	acaatcatca	8760
aacaacaacca	gtcgtatcgt	gtgataatga	gaatatcgca	tgccctgtat	gatgggtctga	8820
gtctagagca	gtctgtgcgc	aaacttcaca	tgctctacaa	cggggagatca	cttttgccac	8880
caacaccaatt	ctcgcggtac	atgcagttata	ctgctgacgg	tcgcgaaagt	ggacatggat	8940
ttttggcgca	gtgtattcaa	aatacgcgca	tgacaataat	gagtgatgac	acgggttggtg	9000
atggaaatcg	tgcaacctgc	aaggcggtgc	acctatcaaa	gattgtcaat	attcctctac	9060
aggtaactcg	aggcagcagt	aacatcatta	ctcaagctac	tgtgtttaac	gcagcctcgcg	9120
cgttagtctt	gtcacgggaa	tctgactcga	aagacgttgt	ctttggcagc	atcgtctctg	9180
gtctgcgaag	cttgcctggt	gaataccagg	acattgtcgg	gcctgtgacc	aacgcagttc	9240
ctgtgtcgcg	tcatatagag	tcgtcagatt	acaaccaatt	gtcgacagtt	atccaagacc	9300
agtcacctct	cagctgtgcca	cacgaaacaa	ttggctctct	agatctcaag	cgcgaactga	9360
cagatgtggc	agaagcaatc	accaactctc	catgctgcat	cacataccac	aatttgcagt	9420
accatccgca	gagtcagttc	gaacagcaga	gagttgagat	gggtgtattg	acaaagtgtg	9480
tcaacattga	gatggatgag	ccactatatg	atttggcgat	tcggggtgaa	gttgaaccag	9540
acggagcagg	actgaaggtt	actgttatcg	cgaagacgca	gttatttggt	aggaagagag	9600
tagaacatct	gtttggaggaa	gtttccaaaa	cgtttgaggg	tctcaactct	ctttgtgaac	9660
gcacgggttg	gtctcaatcg	tcgcgacaga	acaaccgagt	taggtttgta	attcttaatg	9720
acgtttctga	ctttttgggt	tttaccattc	ggagcaataa	gtaagcgaaa	cactggcaaa	9780
tgtcagatat	tacacttcag	aactattatc	ttgactatta	tctcacgttg	tcaggtttca	9840
catgtctgct	acgttgcgat	gagtcacaaa	ttgagatcta	cagggttaacg	caggaaatcca	9900
gaacaattga	caaggattca	tcgatcgaa	actatgattg	gttcgcgtct	ctgacaggac	9960
catctgtcca	ataatagaa	tatagataag	atatcgaggg	aatgcgacaa	ggatgtggaac	10020
ttctcgcgat	agcttgcatt	agtcctctaa	cagtaaaatg	ctcgcgcttc	ttcggattcgt	10080
cttttggtga	tgaattatca	tagtttgttg	tgcgaaagaag	aacgcatgatg	acaggtttgta	10140
ttttggatga	ataaagttc	ataaagttat	gactcttgat	atgataccaa	tatgaaccat	10200
accttgacaa	tattgtgttc	caacattaca	caacacttga	acgggacact	cyttcaacat	10260
caacacaaat	gatctgtcca	acaaagctgc	ctaccttgtc	agtcaccaatg	ggcccaccat	10320
cgaagtcaga	tctgcacctg	tcccaacacc	tggttcagga	gagttgctta	tcaagacaca	10380
tgtctgcgcc	atcaatccag	tcgacggagt	gaagcagctc	atgggttaaca	tgatgtttga	10440
atggctcaag	tatcctctca	tccttggtga	ggcgagggtca	tcaagacggg	tcgaagcggg	10500
acctggtggt	atcgcatatta	aagaaggcga	tagagtttgt	gggtgctacag	caggcatgga	10560
caaggcgagg	agaagtcgcc	acgaaggcgc	atttcaagaa	gtttgcatca	tcgagagagca	10620
tttggtctgt	cgaattccag	agcctgtttac	gtccaccgat	gccacgcttt	tgcccttgac	10680
tttcgcagc	gcgtgcattg	ctttgttcca	aaaggatcaa	ctggcactac	taactcctca	10740
aaccaagatca	aagcgcagtg	caacaagtca	gacagctctg	gtttggggag	cgagtacaag	10800
tggtggggaga	aatgctgtac	agccttgcgt	cgcgggcggc	tatgatgttg	tcgcagacagc	10860
atcacctaag	aactggggata	tcgtacgcgg	tctcgggcgt	tgctcagttt	ttgactatca	10920
cagctcatcg	gccataaacg	atgtggatc	tgctttcaag	gacaagaaat	gcgcagggtgc	10980
tgtagctatt	ggtaagggtg	cactggcgaa	atgcgtcgac	attgtcaaaa	gcgttccggg	11040
agccaccaag	aatgttgcgc	aagttaacct	ctcaatgcct	gagtcacagc	caacaaccaa	11100
gatatactg	attccgtttg	tcgcaaaagta	tttctggatg	gcgggaaactg	atcgactcaa	11160
ggttgcgagc	agtgaggatcc	aaagcaagtt	tggttttggt	acagacataa	tt	11212

<210> 2
 <211> 3129
 <212> PRT
 <213> Fusarium

<400> 2
 Met Glu Tyr Leu Thr Ala Val Asp Gly Arg Gln Asp Leu Pro Pro Thr
 1 5 10 15
 Pro Ala Ser Phe Cys Ser His Gly Asp Ser Pro Leu Asn Ser Ser Tyr
 20 25 30
 Glu Gln Leu Phe His Leu Tyr Gly Leu Asp Ser Ser Arg Ile Glu Ala
 35 40 45
 Ile Lys Pro Cys Thr Pro Phe Gln Leu Asp Met Ile Asp Cys Asn Ala
 50 55 60

Leu Asp Lys Gln Ser Ala Ile Gly His Ala Val Tyr Asp Val Pro Thr
 65 70 75 80
 Asp Ile Asp Ile Ser Arg Phe Ala Leu Ala Trp Lys Glu Ile Val Asn
 85 90 95
 Gln Thr Pro Ala Leu Arg Ala Phe Ala Phe Thr Ser Asp Ser Gly Lys
 100 105 110
 Thr Ser Gln Val Ile Leu Lys Asp Ser Phe Val Phe Ser Trp Met Cys
 115 120 125
 Trp Ser Ser Ser Ser Ser Ser Asp Glu Val Val Arg Asp Glu Ala Ala
 130 135 140
 Ala Ala Ala Ser Gly Pro Arg Cys Asn Arg Phe Val Leu Leu Glu Asp
 145 150 155 160
 Met Gln Thr Lys Lys Cys Gln Leu Val Trp Thr Phe Ser His Ala Leu
 165 170 175
 Val Asp Val Thr Phe Gln Gln Arg Val Leu Ser Arg Val Phe Ala Ala
 180 185 190
 Tyr Lys His Glu Lys Asp Thr His Arg Pro Glu Thr Pro Glu Ser Ser
 195 200 205
 Asp Ala Thr Asp Thr Asp Ser Gln Ser Val Ser Val Val Ser Met Ser
 210 215 220
 Cys Glu Asp Asn Ala Val Ser Ala Thr His Phe Trp Gln Thr His Leu
 225 230 235 240
 Asn Asp Leu Asn Ala Ser Val Phe Pro His Leu Ser Asp His Leu Met
 245 250 255
 Val Pro Asn Pro Thr Thr Thr Ala Glu His Arg Ile Thr Phe Pro Leu
 260 265 270
 Ser Gln Lys Ala Leu Ser Asn Ser Ala Ile Cys Arg Thr Ala Leu Ser
 275 280 285
 Ile Leu Leu Ser Arg Tyr Thr His Ser Asp Glu Ala Leu Phe Gly Ala
 290 295 300
 Val Thr Glu Gln Ser Leu Pro Phe Asp Lys His Tyr Leu Ala Asp Gly
 305 310 315 320
 Thr Tyr Gln Thr Val Ala Pro Leu Arg Val His Cys Gln Ser Asn Leu
 325 330 335
 Arg Ala Ser Asp Val Met Asp Ala Ile Ser Ser Tyr Asp Asp Arg Leu
 340 345 350
 Gly His Leu Ala Pro Phe Gly Leu Arg Asp Ile Arg Asn Thr Gly Asp
 355 360 365
 Asn Gly Ser Ala Ala Cys Asp Phe Gln Thr Val Leu Leu Val Thr Asp
 370 375 380
 Gly Ser His Val Asn Asn Gly Ile Asn Gly Phe Leu Gln Gln Ile Thr
 385 390 395 400
 Glu Ser Ser His Phe Met Pro Cys Asn Asn Arg Ala Leu Leu Leu His
 405 410 415
 Cys Gln Met Glu Ser Ser Gly Ala Leu Leu Val Ala Tyr Tyr Asp His
 420 425 430
 Asn Val Ile Asp Ser Leu Gln Thr Thr Arg Leu Leu Gln Gln Phe Gly
 435 440 445
 His Leu Ile Lys Cys Leu Gln Ser Pro Leu Asp Leu Ser Ser Met Ala
 450 455 460
 Glu Val Asn Leu Met Thr Glu Tyr Asp Arg Ala Glu Ile Glu Ser Trp
 465 470 475 480
 Asn Ser Gln Pro Leu Glu Val Gln Asp Thr Leu Ile His His Glu Met
 485 490 495
 Leu Lys Ala Val Ser His Ser Pro Thr Lys Thr Ala Ile Gln Ala Trp
 500 505 510
 Asp Gly Asp Trp Thr Tyr Ser Glu Leu Asp Asn Val Ser Ser Arg Leu
 515 520 525

Ala Val His Ile Lys Ser Leu Gly Leu Arg Ala Gln Gln Ala Ile Ile
530 535 540
Pro Val Tyr Phe Glu Lys Ser Lys Trp Val Ile Ala Ser Met Leu Ala
545 550 555 560
Val Leu Lys Ser Gly Asn Ala Phe Thr Leu Ile Asp Pro Asn Asp Pro
565 570 575
Pro Ala Arg Thr Ala Gln Val Val Thr Gln Thr Arg Ala Thr Val Ala
580 585 590
Leu Thr Ser Lys Leu His Arg Glu Thr Val Gln Lys Leu Val Gly Arg
595 600 605
Cys Val Val Val Asp Asp Glu Leu Leu Gln Ser Val Ser Ala Ser Asp
610 615 620
Asp Phe Ser Ser Leu Thr Lys Ser Gln Asp Leu Ala Tyr Val Ile Phe
625 630 635 640
Thr Ser Gly Ser Thr Gly Asp Pro Lys Gly Ile Met Ile Glu His Arg
645 650 655
Ala Phe Ser Ser Cys Ala Leu Lys Phe Gly Ala Ser Leu Gly Ile Asn
660 665 670
Ser Asp Thr Arg Ala Leu Gln Phe Gly Thr His Ala Phe Gly Ala Cys
675 680 685
Leu Leu Glu Ile Met Thr Thr Leu Ile Asn Gly Gly Cys Val Cys Ile
690 695 700
Pro Ser Asp Asp Asp Arg Met Asn Ser Ile Pro Ser Phe Ile Asn Arg
705 710 715 720
Tyr Asn Val Asn Trp Met Met Ala Thr Pro Ser Tyr Met Gly Thr Phe
725 730 735
Ser Pro Glu Asp Val Pro Gly Leu Ala Thr Leu Val Leu Val Gly Glu
740 745 750
Gln Met Ser Ser Ser Val Asn Ala Ile Trp Ala Pro Lys Leu Gln Leu
755 760 765
Leu Asn Gly Tyr Gly Gln Ser Glu Ser Ser Ser Ile Cys Phe Ala Ser
770 775 780
Asn Met Ser Thr Glu Pro Asn Asn Met Gly Arg Ala Val Gly Ala His
785 790 795 800
Ser Trp Val Ile Asp Pro Asn Asp Ile Asn Arg Leu Val Pro Ile Gly
805 810 815
Ala Val Gly Glu Leu Val Ile Glu Ser Pro Gly Ile Ala Arg Asp Tyr
820 825 830
Ile Val Pro Pro Pro Glu Lys Ser Pro Phe Phe Thr Asp Ile Pro
835 840 845
Ser Trp Tyr Pro Ala Asn Thr Phe Pro Asp Gly Ala Lys Leu Tyr Arg
850 855 860
Thr Gly Asp Leu Ala Arg Tyr Ala Ser Asp Gly Ser Ile Val Cys Leu
865 870 875 880
Gly Arg Ile Asp Ser Gln Val Lys Ile Arg Gly Gln Arg Val Glu Leu
885 890 895
Gly Ala Ile Glu Thr His Leu Arg Gln Gln Met Pro Asp Asp Leu Thr
900 905 910
Ile Val Val Glu Ala Thr Lys Arg Ser Gln Ser Ala Asn Ser Thr Ser
915 920 925
Leu Ile Ala Phe Leu Ile Gly Ser Ser Tyr Phe Gly Asn Arg Pro Ser
930 935 940
Asp Ala His Ile Leu Asp His Asp Ala Thr Lys Ala Ile Asn Ile Lys
945 950 955 960
Leu Glu Gln Val Leu Pro Arg His Ser Ile Pro Ser Phe Tyr Ile Cys
965 970 975
Met Leu Glu Leu Pro Arg Thr Ala Thr Gly Lys Ile Asp Arg Arg Arg
980 985 990

Leu Arg Ile Met Gly Lys Asp Ile Leu Asp Lys Gln Thr Gln Gly Ala
 995 1000 1005
 Ile Val Gln Gln Ala Pro Ala Pro Ile Pro Val Phe Ala Asp Thr Ala
 1010 1015 1020
 Ala Lys Leu His Ser Ile Trp Val Gln Ser Leu Gly Ile Asp Pro Ala
 1025 1030 1035 1040
 Thr Val Asn Val Gly Ala Thr Phe Phe Glu Leu Gly Gly Asn Ser Ile
 1045 1050 1055
 Thr Ala Ile Lys Met Val Asn Met Ala Arg Ser Val Gly Met Asp Leu
 1060 1065 1070
 Lys Val Ser Asn Ile Tyr Gln His Pro Thr Leu Ala Gly Ile Ser Ala
 1075 1080 1085
 Val Val Lys Gly Asp Pro Leu Ser Tyr Thr Leu Ile Pro Lys Ser Thr
 1090 1095 1100
 His Glu Gly Pro Val Glu Gln Ser Tyr Ser Gln Gly Arg Leu Trp Phe
 1105 1110 1115
 Leu Asp Gln Leu Asp Val Gly Ser Leu Trp Tyr Leu Ile Pro Tyr Ala
 1125 1130 1135
 Val Arg Met Arg Gly Pro Val Asn Val Asp Ala Leu Arg Arg Ala Leu
 1140 1145 1150
 Ala Ala Leu Glu Gln Arg His Glu Thr Leu Arg Thr Thr Phe Glu Asp
 1155 1160 1165
 Gln Asp Gly Val Gly Val Gln Ile Val His Glu Lys Leu Ser Glu Glu
 1170 1175 1180
 Met Lys Val Ile Asp Leu Cys Gly Ser Asp Leu Asp Pro Phe Glu Val
 1185 1190 1195 1200
 Leu Asn Gln Glu Gln Thr Thr Pro Phe Asn Leu Ser Ser Glu Ala Gly
 1205 1210 1215
 Trp Arg Ala Thr Leu Leu Arg Leu Gly Glu Asp Asp His Ile Leu Thr
 1220 1225 1230
 Ile Val Met His His Ile Ile Ser Asp Gly Trp Ser Ile Asp Val Leu
 1235 1240 1245
 Arg Arg Asp Leu Asn Gln Leu Tyr Ser Ala Ala Leu Lys Asp Ser Lys
 1250 1255 1260
 Asp Pro Leu Ser Ala Leu Thr Pro Leu Pro Ile Gln Tyr Ser Asp Phe
 1265 1270 1275 1280
 Ala Lys Trp Gln Lys Asp Gln Phe Ile Glu Gln Glu Lys Gln Leu Asn
 1285 1290 1295
 Tyr Trp Lys Lys Gln Leu Lys Asp Ser Ser Pro Ala Lys Ile Pro Thr
 1300 1305 1310
 Asp Phe Ala Arg Pro Ala Leu Leu Ser Gly Asp Ala Gly Cys Val His
 1315 1320 1325
 Val Thr Ile Asp Gly Glu Leu Tyr Gln Ser Leu Arg Ala Phe Cys Asn
 1330 1335 1340
 Glu His Asn Thr Thr Ser Phe Val Val Leu Leu Ala Ala Phe Arg Ala
 1345 1350 1355 1360
 Ala His Tyr Arg Leu Thr Ala Val Glu Asp Ala Val Ile Gly Thr Pro
 1365 1370 1375
 Ile Ala Asn Arg Asn Arg Pro Glu Leu Glu Asp Ile Ile Gly Cys Phe
 1380 1385 1390
 Val Asn Thr Gln Cys Met Arg Ile Asn Ile Asp His His Asp Thr Phe
 1395 1400 1405
 Gly Thr Leu Ile Asn Gln Val Lys Ala Thr Thr Thr Ala Ala Phe Glu
 1410 1415 1420
 Asn Glu Asp Ile Pro Phe Glu Arg Val Val Ser Ala Leu Gln Pro Gly
 1425 1430 1435 1440
 Ser Arg Asp Leu Ser Ser Thr Pro Leu Ala Gln Leu Ile Phe Ala Val
 1445 1450 1455

His Ser Gln Lys Asp Leu Gly Arg Phe Lys Phe Gln Gly Leu Glu Ser
 1460 1465 1470
 Val Pro Val Pro Ser Lys Ala Tyr Thr Arg Phe Asp Met Glu Phe His
 1475 1480 1485
 Leu Phe Gln Glu Thr Asp Ser Leu Lys Gly Ser Val Asn Phe Ala Asp
 1490 1495 1500
 Glu Leu Phe Lys Met Glu Thr Val Glu Asn Val Val Arg Val Phe Phe
 1505 1510 1515 1520
 Glu Ile Leu Arg Asn Gly Leu Gln Ser Ser Arg Thr Pro Val Ser Ile
 1525 1530 1535
 Leu Pro Leu Thr Asp Gly Ile Val Thr Leu Glu Lys Leu Asp Val Leu
 1540 1545 1550
 Asn Val Lys His Val Asp Tyr Pro Arg Glu Ser Ser Leu Ala Asp Val
 1555 1560 1565
 Phe Gln Thr Gln Val Ser Ala Tyr Pro Asp Ser Leu Ala Val Val Asp
 1570 1575 1580
 Ser Ser Cys Arg Leu Thr Tyr Thr Glu Leu Asp Arg Gln Ser Asp Ile
 1585 1590 1595
 Leu Ala Gly Trp Leu Arg Arg Arg Ser Met Pro Ala Glu Thr Leu Val
 1605 1610 1615
 Ala Val Phe Ala Pro Arg Ser Cys Glu Thr Ile Val Ala Phe Phe Gly
 1620 1625 1630
 Val Leu Lys Ala Asn Leu Ala Tyr Leu Pro Leu Asp Val Arg Ser Pro
 1635 1640 1645
 Ser Ala Arg Val Gln Asp Ile Leu Ser Gly Leu Ser Gly Pro Thr Ile
 1650 1655 1660
 Val Leu Ile Gly His Asp Thr Ala Pro Pro Asp Ile Glu Val Thr Asn
 1665 1670 1675
 Val Glu Phe Val Arg Ile Arg Asp Ala Leu Asn Asp Ser Asn Ala Asp
 1685 1690 1695
 Gly Phe Glu Val Ile Glu His Asp Ser Thr Lys Pro Ser Ala Thr Ser
 1700 1705 1710
 Leu Ala Tyr Val Leu Tyr Thr Ser Gly Ser Thr Gly Arg Pro Lys Gly
 1715 1720 1725
 Val Met Ile Glu His Arg Val Ile Ile Arg Thr Val Thr Ser Gly Cys
 1730 1735 1740
 Ile Pro Asn Tyr Pro Ser Glu Thr Arg Met Ala His Met Ala Thr Ile
 1745 1750 1755
 Ala Phe Asp Gly Ala Ser Tyr Glu Ile Tyr Ser Ala Leu Leu Phe Gly
 1765 1770 1775
 Arg Thr Leu Val Cys Val Asp Tyr Met Thr Thr Leu Asp Ala Arg Ala
 1780 1785 1790
 Leu Lys Asp Val Phe Phe Arg Glu His Val Asn Ala Ala Ser His Val
 1795 1800 1805
 Thr Ser Ser Ser Gln Asp Val Pro Leu Arg Val Pro Arg Arg Leu Ser
 1810 1815 1820
 Arg Thr Leu Met Phe Phe Phe Leu Val Val Thr Asp Ser Thr Ala Pro
 1825 1830 1835
 Asp Ala Leu Asp Ala Gln Gly Leu Tyr Gln Gly Val Gln Cys Tyr Asn
 1845 1850 1855
 Gly Tyr Gly Pro Thr Glu Asn Gly Val Met Ser Thr Ile Tyr Pro Ile
 1860 1865 1870
 Asp Ser Thr Glu Ser Phe Ile Asn Gly Val Pro Ile Gly Arg Ala Leu
 1875 1880 1885
 Asn Asn Ser Gly Ala Tyr Val Val Asp Pro Glu Gln Gln Leu Val Gly
 1890 1895 1900
 Ile Gly Val Met Gly Glu Leu Val Val Thr Gly Asp Gly Leu Ala Arg
 1905 1910 1915 1920

Gly Tyr Ser Asp Lys Ala Leu Asp Glu Asn Arg Phe Val His Ile Thr
 1925 1930 1935
 Val Asn Asp Gln Thr Val Lys Ala Tyr Arg Thr Gly Asp Arg Val Arg
 1940 1945 1950
 Tyr Arg Ile Gly Asp Gly Leu Ile Glu Phe Phe Gly Arg Met Asp Thr
 1955 1960 1965
 Gln Phe Lys Ile Arg Gly Asn Arg Ile Glu Ser Ala Glu Ile Glu Ala
 1970 1975 1980
 Ala Leu Leu Arg Asp Ser Ser Val Arg Asp Ala Ala Val Val Leu Gln
 1985 1990 1995 2000
 Gln Asn Glu Asp Gln Ala Pro Glu Ile Leu Gly Phe Val Val Ala Asp
 2005 2010 2015
 His Asp His Ser Glu Asn Asp Lys Gly Gln Ser Ala Asn Gln Val Glu
 2020 2025 2030
 Gly Trp Gln Asp His Phe Glu Ser Gly Met Tyr Ser Asp Ile Gly Glu
 2035 2040 2045
 Ile Asp Pro Ser Thr Ile Gly Ser Asp Phe Lys Gly Trp Thr Ser Met
 2050 2055 2060
 Tyr Asp Gly Ser Gln Ile Asp Phe Asp Glu Met His Glu Trp Leu Gly
 2065 2070 2075 2080
 Glu Thr Thr Arg Thr Leu His Asp Asn Arg Ser Leu Gly Asn Val Leu
 2085 2090 2095
 Glu Ile Gly Thr Gly Ser Gly Met Ile Leu Phe Asn Leu Asp Ser Arg
 2100 2105 2110
 Leu Glu Ser Tyr Val Gly Leu Glu Pro Ser Arg Ser Ala Ala Ala Phe
 2115 2120 2125
 Val Asn Lys Ala Thr Glu Ser Ile Pro Ser Leu Ala Gly Lys Ala Lys
 2130 2135 2140
 Val Gln Val Gly Thr Ala Thr Asp Ile Gly Gln Val Asp Asp Leu His
 2145 2150 2155 2160
 Pro Asp Leu Val Val Leu Asn Ser Val Ile Gln Tyr Phe Pro Ser Ser
 2165 2170 2175
 Glu Tyr Leu Ala Glu Ile Ala Asp Thr Leu Ile His Leu Pro Asn Val
 2180 2185 2190
 Gln Arg Ile Phe Phe Gly Asp Val Arg Ser Gln Ala Thr Asn Glu His
 2195 2200 2205
 Phe Leu Ala Ala Arg Ala Ile His Thr Leu Gly Lys Asn Ala Thr Lys
 2210 2215 2220
 Asp Asp Val Arg Gln Lys Met Ala Glu Leu Glu Asp Met Glu Glu Glu
 2225 2230 2235 2240
 Leu Leu Val Glu Pro Ala Phe Phe Thr Ser Leu Lys Asp Arg Phe Pro
 2245 2250 2255
 Gly Leu Val Glu His Val Glu Ile Leu Pro Lys Asn Met Glu Ala Val
 2260 2265 2270
 Asn Glu Leu Ser Ala Tyr Arg Tyr Ala Ala Val Val His Val Arg Gly
 2275 2280 2285
 Ser Leu Gly Asp Glu Leu Val Leu Pro Val Glu Lys Asp Asp Trp Ile
 2290 2295 2300
 Asp Phe Gln Ala Asn Gln Leu Asn Gln Lys Ser Leu Gly Asp Leu Leu
 2305 2310 2315 2320
 Lys Ser Ser Asp Ala Ala Ile Met Ala Val Ser Lys Ile Pro Phe Glu
 2325 2330 2335
 Ile Thr Ala Phe Glu Arg Gln Val Val Ala Ser Leu Asn Ser Asn Ile
 2340 2345 2350
 Asp Glu Trp Gln Leu Ser Thr Ile Arg Ser Ser Ala Glu Gly Asp Ser
 2355 2360 2365
 Ser Leu Ser Val Pro Asp Ile Phe Arg Ile Ala Gly Glu Ala Gly Phe
 2370 2375 2380

Arg Val Glu Val Ser Ser Ala Arg Gln Trp Ser Gln Asn Gly Ala Leu
 2385 2390 2395 2400
 Asp Ala Val Phe His His Cys Cys Ser Gln Gly Arg Thr Leu Val Asn
 2405 2410 2415
 Phe Pro Thr Asp His His Leu Arg Gly Ser Asp Leu Leu Thr Asn Arg
 2420 2425 2430
 Pro Leu Gln Arg Leu Gln Asn Arg Arg Ile Ala Ile Glu Val Arg Glu
 2435 2440 2445
 Arg Leu Arg Ser Leu Leu Pro Ser Tyr Met Ile Pro Ser Asn Ile Val
 2450 2455 2460
 Val Leu Asp Lys Met Pro Leu Asn Ala Asn Gly Lys Val Asp Arg Lys
 2465 2470 2475 2480
 Glu Leu Ser Arg Arg Ala Lys Val Val Pro Lys Gln Gln Thr Ala Ala
 2485 2490 2495
 Pro Leu Pro Thr Phe Pro Ile Ser Glu Val Glu Val Ile Leu Cys Glu
 2500 2505 2510
 Glu Ala Thr Glu Val Phe Gly Met Lys Val Asp Ile Thr Asp His Phe
 2515 2520 2525
 Phe Asn Leu Gly Gly His Ser Leu Leu Ala Thr Lys Leu Ile Ser Arg
 2530 2535 2540
 Ile Asp Gln Arg Leu Lys Val Arg Ile Thr Val Lys Asp Val Phe Asp
 2545 2550 2555 2560
 His Pro Val Phe Ala Asp Leu Ala Ser Val Ile Arg Gln Gly Leu Gly
 2565 2570 2575
 Leu Gln Gln Pro Val Ser Asp Gly Gln Gly Gln Asp Arg Ser Ala His
 2580 2585 2590
 Met Ala Pro Arg Thr Glu Thr Glu Ala Ile Leu Cys Asp Glu Phe Ala
 2595 2600 2605
 Lys Val Leu Gly Phe Gln Val Gly Ile Thr Asp Asn Phe Phe Asp Leu
 2610 2615 2620
 Gly Gly His Ser Leu Met Ala Thr Lys Leu Ala Val Arg Ile Gly His
 2625 2630 2635 2640
 Arg Leu Asp Thr Thr Val Ser Val Lys Asp Val Phe Asp His Pro Val
 2645 2650 2655
 Leu Phe Gln Leu Ala Ile Ala Leu Asp Asn Leu Val Gln Ser Lys Thr
 2660 2665 2670
 Asn Glu Ile Val Gly Gly Arg Glu Met Ala Glu Tyr Ser Pro Phe Gln
 2675 2680 2685
 Leu Leu Phe Thr Glu Asp Pro Glu Glu Phe Met Ala Ser Glu Ile Lys
 2690 2695 2700
 Pro Gln Leu Glu Leu Gln Glu Ile Ile Gln Asp Ile Tyr Pro Ser Thr
 2705 2710 2715 2720
 Gln Met Gln Lys Ala Phe Leu Phe Asp His Thr Thr Ala Arg Pro Arg
 2725 2730 2735
 Pro Phe Val Pro Phe Tyr Ile Asp Phe Pro Ser Thr Ser Glu Pro Asp
 2740 2745 2750
 Ala Ala Gly Leu Ile Lys Ala Cys Glu Ser Leu Val Asn His Leu Asp
 2755 2760 2765
 Ile Phe Arg Thr Val Phe Ala Glu Ala Ser Gly Glu Leu Tyr Gln Val
 2770 2775 2780
 Val Leu Ser Cys Leu Asp Leu Pro Ile Gln Val Ile Glu Thr Glu Asp
 2785 2790 2795 2800
 Asn Ile Asn Thr Ala Thr Asn Glu Phe Leu Asp Glu Phe Ala Lys Glu
 2805 2810 2815
 Pro Val Arg Leu Gly His Pro Leu Ile Arg Phe Thr Ile Ile Lys Gln
 2820 2825 2830
 Thr Lys Ser Met Arg Val Ile Met Arg Ile Ser His Ala Leu Tyr Asp
 2835 2840 2845

Gly Leu Ser Leu Glu His Val Val Arg Lys Leu His Met Leu Tyr Asn
 2850 2855 2860
 Gly Arg Ser Leu Leu Pro Pro His Gln Phe Ser Arg Tyr Met Gln Tyr
 2865 2870 2875 2880
 Thr Ala Asp Gly Arg Glu Ser Gly His Gly Phe Trp Arg Asp Val Ile
 2885 2890 2895
 Gln Asn Thr Pro Met Thr Ile Leu Ser Asp Asp Thr Val Val Asp Gly
 2900 2905 2910
 Asn Asp Ala Thr Cys Lys Ala Leu His Leu Ser Lys Ile Val Asn Ile
 2915 2920 2925
 Pro Ser Gln Val Leu Arg Gly Ser Ser Asn Ile Ile Thr Gln Ala Thr
 2930 2935 2940
 Val Phe Asn Ala Ala Cys Ala Leu Val Leu Ser Arg Glu Ser Asp Ser
 2945 2950 2955 2960
 Lys Asp Val Val Phe Gly Arg Ile Val Ser Gly Arg Gln Gly Leu Pro
 2965 2970 2975
 Val Glu Tyr Gln Asp Ile Val Gly Pro Cys Thr Asn Ala Val Pro Val
 2980 2985 2990
 Arg Ala His Ile Glu Ser Ser Asp Tyr Asn Gln Leu Leu His Asp Ile
 2995 3000 3005
 Gln Asp Gln Tyr Leu Leu Ser Leu Pro His Glu Thr Ile Gly Phe Ser
 3010 3015 3020
 Asp Leu Lys Arg Asn Cys Thr Asp Trp Pro Glu Ala Ile Thr Asn Phe
 3025 3030 3035 3040
 Ser Cys Cys Ile Thr Tyr His Asn Phe Glu Tyr His Pro Glu Ser Gln
 3045 3050 3055
 Phe Glu Gln Gln Arg Val Glu Met Gly Val Leu Thr Lys Phe Val Asn
 3060 3065 3070
 Ile Glu Met Asp Glu Pro Leu Tyr Asp Leu Ala Ile Ala Gly Glu Val
 3075 3080 3085
 Glu Pro Asp Gly Ala Gly Leu Lys Val Thr Val Ile Ala Lys Thr Gln
 3090 3095 3100
 Leu Phe Gly Arg Lys Arg Val Glu His Leu Leu Glu Glu Val Ser Lys
 3105 3110 3115 3120
 Thr Phe Glu Gly Leu Asn Ser Ser Leu
 3125

<210> 3
 <211> 23
 <212> DNA
 <213> Fusarium

<400> 3
 aattgattcg cttgaaagtc gat

23

<210> 4
 <211> 25
 <212> DNA
 <213> Fusarium

<400> 4
 cttgagagtt acgttggtct tgaac

25